

## New research reveals how grazing management practices affect cattle weight gain by altering foraging behavior

May 9 2023, by Maribel Alonso



Cattle moving in the pasture. Credit: USDA-ARS Rangeland Resources & Systems Research

Rotational or continuous grazing? Which system allows for more sustainable and profitable free-range livestock production?

Ranchers often rotate cattle seasonally among different pastures on their



operations but may allow cattle to graze season-long in a single pasture. Implementing a more intensive rotational system within the growing season—dividing the pasture into smaller areas, or paddocks, and shifting a herd of animals throughout the season—has been suggested to offer a greater chance for more sustainable grazing management. However, there have been very few experimental studies testing these ideas in extensive grazing systems.

A team of researchers at the USDA's Agricultural Research Service is completing a 10-year study on how grazing practices used in these two systems affect cattle foraging behavior, diet quality, and yearly <u>weight</u> <u>gain</u> in semi-arid, extensive rangelands.

The team started by exploring the efficiency of using cattle <u>global</u> <u>positioning system</u> (GPS) tracking collars, combined with activity sensors, to monitor the animals' grazing activities.

"The primary objective of the study was using the sensors to measure the steers' foraging behavior, and relate this to how they are gaining weight," said David Augustine, a research ecologist with the ARS Rangeland Resources & Systems Research in Colorado.

The collars collected precise data based on the animals' feeding habits per day, such as how much time was spent grazing every day, how many steps were taken (grazing speed), the shape of the foraging pathways, and how long each animal lowered their head as a sign of eating (representing the length of meals).

Even though the accuracy of the application is still being refined, results showed that the technology can inform livestock managers about animal distribution and foraging behaviors of free-ranging cattle in extensive rangelands. Being able to monitor foraging <u>behavior</u> continuously in turn allows managers to make more timely decisions on how, when, and



where to move cattle within their operation, or to sell cattle at optimal times.

The researchers then applied the same technology in a range-scale experiment where steers were either divided into smaller herds in the paddocks of a non-rotational (season-long) grazing system or managed as a single large herd in a multi-paddock rotational system (using a collaborative and adaptive rangeland management approach). The first five years of data showed that the rotationally managed cattle gained, on average, 14% less weight than <u>cattle</u> in the season-long management system.

"Cattle did not have as much freedom to roam and be more selective on what to eat, so they ended up eating what was available in front of them, which was <u>lower quality</u> forages with less protein, and they gained less weight as a result. These behaviors were directly correlated to the average of 14% reduction in steer weight gain in the multi-paddock rotating system compared to the non-rotational continuous system," explained Augustine.

The study showed herds in the multi-paddock rotating system feeding in more linear pathways instead of moving around looking for greener grass and selecting bites of more digestible vegetation. They also fed slower, spent more time on the same patch of grass, and didn't turn their heads around much while feeding, compared with steers in the continuous grazing system. These behaviors of less selective <u>foraging</u> resulted in a lower diet quality, which led to reduced weight gain during the growing season.

Results, published in *Agriculture, Ecosystems, and Environment*, show that large herds grazing in small, homogenous paddocks have little opportunity to move around in ways that let them feed on high quality diet.



**More information:** David J. Augustine et al, Adaptive, multi-paddock, rotational grazing management alters foraging behavior and spatial grazing distribution of free-ranging cattle, *Agriculture, Ecosystems & Environment* (2023). DOI: 10.1016/j.agee.2023.108521

## Provided by Agricultural Research Service

Citation: New research reveals how grazing management practices affect cattle weight gain by altering foraging behavior (2023, May 9) retrieved 1 May 2024 from <u>https://phys.org/news/2023-05-reveals-grazing-affect-cattle-weight.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.