

Study reveals presence of Hunga Tonga eruption aerosols in Northern Hemisphere stratospheric westerlies

May 5 2023, by Li Yuan



Dr. Bai Zhixuan and his colleagues launched a balloon in Golmud, Qinghai province on March 2021, as part of the SWOP campaign. Credit: Chen Zhixiong

An international team led by researchers from the Institute of Atmospheric Physics (IAP) of the Chinese Academy of Sciences has detected aerosols from the Hunga Tonga eruption in the Northern



Hemisphere stratospheric westerlies for the first time.

This discovery, published in *Science Bulletin*, will aid in simulating the physical and chemical processes in <u>volcanic plumes</u>.

The Hunga Tonga-Hunga Ha'apai volcano, located in the South Pacific Ocean, erupted in January 2022, spewing ash and gas into the atmosphere. The eruption was one of the largest in recent years and was closely monitored by scientists worldwide.

The purpose of the Sounding Water vapor, Ozone, and Particle (SWOP) campaign is to better understand the temporal and spatial distribution characteristics of atmospheric compositions over the Tibetan Plateau. Capturing the <u>volcanic aerosols</u> from the Hunga Tonga eruption was unexpected, according to Prof. Bian Jianchun, an IAP scientist and a team leader of the SWOP campaign.

"We happen to have captured the volcanic <u>aerosols</u> from the Hunga Tonga eruption 12 weeks after it erupted. To my knowledge, this is the first time that it's been detected at the stratospheric westerlies in the Northern Hemisphere," said Prof. Bian.

Large uncertainties still remain in understanding physical properties and evolutionary process of volcanic aerosols in the stratosphere, particularly SO_2 oxidation in volcanic plumes and clouds and the interaction between ash and sulfate aerosols after volcanic eruptions. In-situ measurements obtained in volcanic plumes, especially measurements of the microphysical parameters of volcanic aerosols, provide compelling evidence that reduces these uncertainties. However, it is difficult to obtain these parameters, especially during the occurrence of volcanic eruptions.

During the SWOP campaign carried out over the Tibetan Plateau, a



balloon-borne payload consisting of the Compact Optical Backscatter Aerosol Detector (COBALD) and the Portable Optical Particle Spectrometer (POPS) was released in Lijiang, Yunan province on April 9, 2022. Unexpectedly, the balloon captured the size spectrum of volcanic aerosols at 24–25 km.

The Hunga Tonga volcanic aerosol was identified by the size distribution of the aerosol diameter, which was centered at approximately 0.42-1.27 µm, larger than the background mode (0.22-0.42 µm) of stratospheric aerosols, and particle number densities of approximately 1 cm⁻³. The backscatter ratios derived from the COBALD measurements at 455 nm and 940 nm both showed enhanced aerosol signals between 24 and 25 km. Additionally, measurements showed that the aerosol and water vapor layers were separated in the Hunga Tonga plume.

These findings suggest that <u>volcanic eruptions</u> have potential impact on regional and global atmospheric circulation and climate, even thousands of kilometers away from the source. The possible mechanisms that led to the separation of aerosol and water vapor layers in the Hunga Tonga plume will be examined in future study.

More information: Jianchun Bian et al, First detection of aerosols of the Hunga Tonga eruption in the Northern Hemisphere stratospheric westerlies, *Science Bulletin* (2023). DOI: 10.1016/j.scib.2023.03.002

Provided by Chinese Academy of Sciences

Citation: Study reveals presence of Hunga Tonga eruption aerosols in Northern Hemisphere stratospheric westerlies (2023, May 5) retrieved 27 June 2024 from <u>https://phys.org/news/2023-05-reveals-presence-hunga-tonga-eruption.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.