

## How plants use sugar to produce roots

## May 22 2023



Two lateral root primordia develop from the main root of Arabidopsis thaliana.



The images (with false colors) were taken with a confocal microscope. Credit: Michael Stitz, Heidelberg University

Along with sugar reallocation, a basic molecular mechanism within plants controls the formation of new lateral roots. An international team of plant biologists has demonstrated that it is based on the activity of a certain factor, the target of rapamycin (TOR) protein. A better understanding of the processes that regulate root branching at the molecular level could contribute to improving plant growth and therefore crop yields, according to research team leader Prof. Dr. Alexis Maizel of the Centre for Organismal Studies at Heidelberg University.

Good root growth ensures that <u>plants</u> can absorb sufficient nutrients and grow, thus contributing to their general fitness. To do that, they must align the available resources from <u>metabolic processes</u> with their genetic development programs. Plants bind <u>carbon dioxide</u> (CO<sub>2</sub>) from the atmosphere in their leaves and convert it to simple sugars via photosynthesis. In the form of fructose and glucose, these simple sugars are also allocated in the roots, where they drive the growth and development of the plant.

Prof. Maizel's team used the thale cress Arabidopsis thaliana, a model plant in plant research, to study how this process occurs at the molecular level. Their investigations focus on what role glucose plays in forming lateral roots. "We do know that, besides plant hormones, sugar from the shoot is also allocated in the roots, but how the plant recognizes that sugar resources are available for forming lateral roots has not been understood thus far," explains Dr. Michael Stitz, a researcher on Alexis Maizel's team.

The studies at the metabolism level showed that Arabidopsis forms



lateral roots only when glucose breaks down and carbohydrates are consumed in the pericycle—the outermost cell layer of the main root cylinder. This process is controlled at the <u>molecular level</u> by the target of rapamycin protein. This factor controls critical signal networks and metabolic processes in plants as well as in animals and humans. Its activity is governed by the interaction of growth factors like the plant hormone auxin and nutrients like sugar.

Using Arabidopsis, the researchers discovered that TOR becomes active in the pericycle cells only when sugar is present there. So-called founder cells then form the lateral roots through cell division.

Prof. Maizel says, "TOR assumes a kind of gatekeeper role; when the plant activates the genetic growth program responsible for <u>root</u> formation via the hormone auxin, TOR checks whether there are sufficient sugar resources available for this process."

TOR acts by controlling the translation of specific auxin-dependent genes, blocking their expression if there aren't sufficient <u>sugar</u> resources available. When the researchers suppressed TOR activity, no lateral roots were formed. "That suggests that a fundamental molecular mechanism is involved," states the Heidelberg plant biologist.

At the same time, the researchers demonstrated that TOR controls, via a similar mechanism, the formation of roots from other <u>plant tissues</u>—the so-called adventitious roots. According to Prof. Maizel, the results from their investigations could also be of interest for agricultural applications. "They could potentially be used to develop new strategies for <u>plant</u> growth optimized for various environmental conditions and better crop yields," says the researcher.

The findings are published in The EMBO Journal.



**More information:** Michael Stitz et al, TOR acts as a metabolic gatekeeper for auxin-dependent lateral root initiation in Arabidopsis thaliana, *The EMBO Journal* (2023). DOI: 10.15252/embj.2022111273

## Provided by Heidelberg University

Citation: How plants use sugar to produce roots (2023, May 22) retrieved 28 June 2024 from <u>https://phys.org/news/2023-05-sugar-roots.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.