

Zeroing in on a fundamental property of the proton's internal dynamics

May 1 2023



A proton with transverse spin and quarks inside also with transverse spins. The tensor charge can be calculated for "up" and "down" quarks by various methods to quantify their total transverse spin in the proton. Credit: Thomas Jefferson National Accelerator Facility



Inside the proton are elementary particles called quarks. Quarks and protons have an intrinsic angular momentum called spin. Spin can point in different directions. When it is perpendicular to the proton's momentum, it is called a transverse spin. Just like the proton carries an electric charge, it also has another fundamental charge called the tensor charge. The tensor charge is the net transverse spin of quarks in a proton with transverse spin.

The only way to obtain the <u>tensor</u> charge from <u>experimental data</u> is using the theory of quantum chromodynamics (QCD) to extract the "transversity" function. This universal function encodes the difference between the number of quarks with their spin aligned and anti-aligned to the proton's spin when it is in a transverse direction. Using state-of-theart data science techniques, researchers recently made the most precise empirical determination of the tensor charge.

Due to the phenomenon known as confinement, quarks are always bound in the proton or other hadrons (particles with multiple quarks). The challenge is to connect the theory of <u>quark</u> interactions (QCD) to experimental measurements of high-energy collisions involving hadrons.

Researchers from the Coordinated Theoretical Approach to Transverse Momentum Dependent Hadron Structure in QCD Topical Collaboration (TMD Collaboration), working in conjunction with the Thomas Jefferson National Accelerator Facility (Jefferson Lab) Angular Momentum Collaboration (JAM Collaboration), analyzed data from a wide range of experiments where protons and/or quarks were transversely polarized. This allowed for the most precise empirical determination of the proton's tensor charge. The tensor charge is not only a fundamental property of the proton but also needed in searches for new physics.

The results were then compared to computations of the proton's tensor



charge by lattice QCD, which simulates the proton's structure on a supercomputer. After about a decade of results showing disagreement between empirical methods and lattice QCD for the proton's tensor charge, researchers for the first time found agreement between the two. Their findings are published in the journal *Physical Review D*.

The <u>empirical study</u> was performed using QCD theory and state-of-theart numerical methods. A crucial part of the analysis was the utilization of data from electron–positron, electron–proton, and proton–proton scattering. This opens a new frontier in QCD global analyses to simultaneously include all possible measurements, like those from the future Electron-Ion Collider and Jefferson Lab 12 GeV, to continue to increase the precision and accuracy of extracting the proton's tensor charge.

More information: Leonard Gamberg et al, Updated QCD global analysis of single transverse-spin asymmetries: Extracting H^{\sim} , and the role of the Soffer bound and lattice QCD, *Physical Review D* (2022). DOI: 10.1103/PhysRevD.106.034014

Provided by US Department of Energy

Citation: Zeroing in on a fundamental property of the proton's internal dynamics (2023, May 1) retrieved 23 June 2024 from <u>https://phys.org/news/2023-05-zeroing-fundamental-property-proton-internal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.