

Chemical communication between female rats exists and is complex

June 28 2023



Protein profile of female rat scent marks. Female rat scent marks were recovered by swabbing and analyzed by SDS-PAGE from Wistar Han (WH), wild rats (Wild), and Brown Norway (BN). The major bands were excised and analyzed by PMF; band identities are indicated on the right-hand side of each gel (A). Female rat scent marks were analyzed by label-free quantitative proteomics. The quantitative profile, averaged across all samples (with error bars denoting



SEM), covered six orders of magnitude. Key proteins: MUPs, beta glucuronidase, and protease/antiprotease protein are highlighted (B). The proteins in scent marks were also analyzed by electrospray ionization mass spectrometry to recover intact protein masses, revealing to a family of masses around 17 kDa, readily reconciled to partial trimming of MUP1 (C). The sites of trimming are readily accessible in a flexible unstructured region at the N terminus [D, based on a predicted model of MUP1; built using the Phyre]. Credit: *Proceedings of the National Academy of Sciences* (2023). DOI: 10.1073/pnas.2300794120

Scent marking is a communication strategy for many mammals. These scent marks provide key information about the animals that leave them, with those animals that detect them often changing their behavior in response to them.

This <u>chemical communication</u> has been studied in depth when it comes to males, but, in the case of <u>females</u>, scant attention had been paid to how they use it, to be receptive to males, or to communicate with their offspring. This dynamic had given rise to a <u>knowledge gap</u> regarding <u>communication</u> between females, which, in <u>social species</u> where they have to share territory, is fundamental, not only in terms of competition, but also cooperation.

A clear example of this shortcoming in knowledge was that of <u>female</u> <u>rats</u>, which are a species often studied to understand animals' cognitive and psychological behavior. A new study published in the journal *PNAS*, in which University of Cordoba researcher Guadalupe Gómez Baena worked together with a team from the University of Liverpool, demonstrates that female rats communicate with each other through <u>scent marks</u> deposited in their environments.

The direct study of the marks left by populations of Norway rats (Rattus



norvegicus) demonstrates that the rats use different, complex signals to communicate with each other. "It tends to be thought that rats are very similar to mice, and that the two species use the same method to communicate. Our article shows that the communication strategy is completely different in these species. Moreover, females have never received much attention, so our article tries to shed light on communication between females," explained the researcher with the UCO's Department of Biochemistry and Molecular Biology.

At this point, MUP (major urinary proteins) enter into play. MUPs are key in the transmission of odors through marks, yet function differently in mice (found in the urine with which they mark) as opposed to rats (whose marks contains sebaceous secretions, in addition to urine).

"In this work we analyze, for the first time, the marks that the animals deposit. Finding that in females they contain a wide variety of proteins from the clitoral glands, including a shorter version of MUPs," says Gómez Baena. This article demonstrates the connection between the clitoral glands responsible for these sebaceous secretions, urine, and the marks left by rats. Thus proving that these proteins, shorter in females, come from the clitoral glands. Therefore, the combination of urine and the secretion from the clitoral glands is necessary for the marks of the females to arouse interest in other females.

What do females 'talk about?'

When and why do female interrelate? To find out whether females use marks differently—depending on whether they are in heat, or the genetic identity of their counterparts—behavioral tests were carried out to determine the <u>social context</u> in which the females communicate.

Regardless of whether they are in heat, or the stage, females communicate with others by marking territory, because those marks may



not be only to compete, but to cooperate as well. In terms of competition, they also use that communication to verify whether their counterparts are in heat, which may be beneficial during mating periods.

When responding to different marks, genetic background also matters: females increase their marking in response to that of other females of the same breed, which could reflect a motivation to communicate with females of the same colony.

This study focuses on communication between female rats, expanding our knowledge of a species widely used in experimentation and demonstrating the need to study the specificities of the females of other species.

More information: Guadalupe Gómez-Baena et al, Unraveling female communication through scent marks in the Norway rat, *Proceedings of the National Academy of Sciences* (2023). DOI: 10.1073/pnas.2300794120

Provided by University of Córdoba

Citation: Chemical communication between female rats exists and is complex (2023, June 28) retrieved 10 May 2024 from <u>https://phys.org/news/2023-06-chemical-communication-female-rats-complex.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.