


# Effect of nitrogen deposition on animal species stronger than expected

June 12 2023



Conceptual representation of the fitness responses to changed producer tissue N:P ratio of a generalist versus specialist consumer (A), and species that differ in the location of the threshold elemental ratio (TER) due to different life-history strategies, here a fast-growing versus slow-growing consumer (B). Peaks of the quadratic curves represent the TER; fitness loss occurs when deviating from this TER. The solid vertical line denotes producer N:P ratio under ambient conditions, and the dashed vertical line shows increased producer N:P ratios as result of N deposition (N dep.), indicated by the horizontal arrow. Solid horizontal lines indicate fitness of the corresponding consumer under ambient conditions, and dashed horizontal lines show the resulting fitness under N-deposition-mediated increases in N:P ratio. Vertical arrows indicate the degree and direction of fitness shifts resulting from a changed plant tissue N:P ratio. Credit: *Biological Reviews* (2023). DOI: 10.1111/brv.12972

Anthropogenic increased nitrogen deposition is a well-known environmental stressor, resulting in impoverishment of soil quality in naturally nutrient-poor ecosystems. As a result, habitat conditions for plant and animal species are also changing. Such changes are sometimes clearly visible in the field. For instance, plant species that benefit from extra nitrogen often become dominant, leading to grass-dominance in heathlands and increasing cover of blueberries and bramble in woodlands. Less visible however are the effects of nitrogen deposition on food quality of plants, and the subsequent effects on animal fitness.

A group of researchers from several Dutch and Belgian universities and research institutes show that [nitrogen deposition](#) alters the ratio of nutrients in plants, which in turn may lead to a reduced performance in many [animal species](#). The findings were recently published in the journal *Biological Reviews*.

The ratio of nitrogen with other important elements such as phosphorus and calcium shifts, and plants often produce more nitrogen-rich amino acids relative to other [essential amino acids](#) under increased nitrogen load. As a result, shortages of necessary nutrients for animals can occur more often. Plants also invest less in carbon-rich defense chemicals, while nitrogen-rich toxins increase.

Animal species that grow rapidly and insects that perform a complete metamorphosis, from larva to adult via a pupal stage, such as butterflies, bees and flies, require relatively high levels of phosphorus in order to sustain this high growth speed. These species are thus more affected by changes in plant quality than slow-growing species and insects that develop as nymphs, such as bugs and grasshoppers.

By comparing these and other traits of species, the researchers

concluded that a small group of generalist species likely benefits from the extra nitrogen and pest outbreaks will occur more often, but a much larger group of animal [species](#) suffers from changed nutrient content and thus will decline or disappear under increased nitrogen load.

**More information:** Joost J. Vogels et al, Towards a mechanistic understanding of the impacts of nitrogen deposition on producer–consumer interactions, *Biological Reviews* (2023). [DOI: 10.1111/brv.12972](#)

Provided by Radboud University

Citation: Effect of nitrogen deposition on animal species stronger than expected (2023, June 12) retrieved 2 May 2024 from <https://phys.org/news/2023-06-effect-nitrogen-deposition-animal-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.