


Improving fluid simulations with embedded neural networks

June 2 2023, by Samuel Jarman



Simulating flows in a complex fluid. Credit: *The European Physical Journal E* (2023). DOI: 10.1140/epje/s10189-023-00267-w

While neural networks can help to improve the accuracy of fluid flow simulations, new research shows how their accuracy is limited unless the right approach is taken. By embedding fluid properties into neural networks, simulation accuracy can improve by orders of magnitude.

The Lattice Boltzmann Method (LBM) is a simulation technique used to

describe the dynamics of fluids. Recently, there has been an increasing interest in employing neural networks for computational modeling of fluids. The results of a collaboration between researchers from Eindhoven University of Technology and Los Alamos National Laboratory, published in *The European Physical Journal E*, show how [neural networks](#) can be embedded into a LBM framework to model collisions between fluid particles.

The team found that it is essential to embed the correct physical properties into the neural network architecture to preserve accuracy. These discoveries could deepen researchers' understanding of how to model fluid flows.

The LBM works by dividing flows into grids of tiny cells: tracking the movement of particles between cells, then calculating how the distribution of particles in each cell changes over time. One of the LBM's key requirements is to model collisions between particles. Several different models are now capable of doing this, and by appropriately recreating these collisions within the LBM, researchers can now use them to model a diverse array of flow systems.

In their study, the Eindhoven and Los Alamos teams assessed the reliability of this approach by using the LBM to model time-varying behavior in a variety of complex flows. They discovered that 'vanilla' neural network architectures, with no enforcement of physical properties, have a very limited accuracy when calculating particle collisions.

Yet by embedding the unique physical properties of real flows—including conservation laws and spatial symmetries—their [accuracy](#) in reproducing time-varying dynamics can improve by some 4 orders of magnitude. The authors hope their discoveries could have important implications for [fluid dynamics](#): allowing researchers to

simulate systems including mixtures of gases, turbulent flows, as well as more exotic quantum fluids.

More information: Alessandro Corbetta et al, Toward learning Lattice Boltzmann collision operators, *The European Physical Journal E* (2023).

[DOI: 10.1140/epje/s10189-023-00267-w](https://doi.org/10.1140/epje/s10189-023-00267-w)

Provided by Springer

Citation: Improving fluid simulations with embedded neural networks (2023, June 2) retrieved 23 June 2024 from

<https://phys.org/news/2023-06-fluid-simulations-embedded-neural-networks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.