


Plant functional trait diversity increases soil organic carbon sequestration during tropical forest succession

June 15 2023, by Zhang Nannan



Credit: Unsplash/CC0 Public Domain

Previous studies have shown that soil organic carbon (SOC) influences the provision of edaphic ecosystem services, making it a key indicator of

soil health under sustainable land use management and for climate change resilience. However, the SOC sequestration processes during tropical forest succession remain unclear, particularly after land abandonment from agriculture practices.

In a study published in the *Journal of Ecology*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) and the Kunming Institute of Botany of the Chinese Academy of Sciences have investigated the effects of plant [functional diversity](#), species diversity, phylogenetic diversity, tree above-ground biomass, and [environmental factors](#) on SOC during forest succession following land abandonment from swidden agriculture.

The researchers used data from two large-scale surveys of 320 plots (of 1,000 m² each) and 1,280 subplots (of 100 m² each) covering a total landscape area of 20,000 ha in Southeast Asia. They measured SOC stocks during forest succession to infer sequestration via space-time substitution.

They found that functional trait diversity played an important role in determining SOC sequestration across successional trajectories. Plant trait composition was directly associated with plants-to-soil carbon inputs and outputs, reflecting complementary resource use within [plant communities](#) and their impact on biogeochemical cycles.

In addition, increases in SOC carbon storage were associated with indirect positive effects of [species diversity](#) and succession age via functional trait diversity, but phylogenetic diversity and aboveground biomass showed no [significant relationship](#) with SOC stocks.

Furthermore, the effects of soil properties and functional trait diversity on SOC carbon storage shifted with elevation.

"Our study helps to clarify the relationships between different components of plant biodiversity, environmental factors, and soil carbon stocks in natural tropical forest succession, especially after land abandonment by swidden agriculture," said Gbadamassi Dossa, corresponding author of the study.

The researchers suggested that in the mosaic landscape of the tropical region, a trait-based approach in restoration and afforestation management practices should be considered.

More information: Manichanh Satdichanh et al, Drivers of soil organic carbon stock during tropical forest succession, *Journal of Ecology* (2023). [DOI: 10.1111/1365-2745.14141](https://doi.org/10.1111/1365-2745.14141)

Provided by Chinese Academy of Sciences

Citation: Plant functional trait diversity increases soil organic carbon sequestration during tropical forest succession (2023, June 15) retrieved 2 May 2024 from <https://phys.org/news/2023-06-functional-trait-diversity-soil-carbon.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|