


Protective molecules facilitate molybdate anion binding to create novel silver nanoclusters

June 20 2023



The silver-molybdate nanocluster is supported by thiolate and phosphine molecules, which serve as protective ligands that surround the silver nanocluster to increase stability and uniformity. Silver atoms are blue, molybdate ions are orange and red, thiolate ligands are yellow and gray and phosphine ligands are purple and gray. Credit: *Polyoxometalates*, Tsinghua University Press

Silver nanocluster synthesis, or the binding of a silver atom to negatively charged anions, is often plagued by stability, sizing uniformity and surface composition issues. Protective molecules that bind the surface of silver nanoclusters, called ligands, can mitigate these issues, creating new nanoclusters with unique physical and chemical properties.

Nanoclusters are stable, crystalline materials up to two nanometers in diameter that are engineered for specific properties, such as the acceleration of a particular chemical reaction, as a semiconducting material or for targeted drug delivery.

A group of nanoparticle scientists from Mian-Ming Zhang's group at Shanxi Normal University and the Taiyuan University of Technology recently synthesized and characterized a new silver-molybdate nanocluster with the help of thiolate and phosphine [ligands](#) to protect the surface of the [cluster](#) and increase stability and uniformity. The use of protective ligands will assist other researchers in synthesizing additional silver metal and molybdate anion nanoclusters with unique characteristics in the future.

The team published the results of their study in the May 11 issue of the journal *Polyoxometalates*.

"In order to further stabilize the cluster during synthesis, or to obtain a diverse cluster structure, a second ligand or a larger variety of ligands is often introduced during nanocluster synthesis. Strategies for mixed ligand protection were chosen in our study. For thiol/phosphine mixed ligands, adding thiol can overcome the single coordination of phosphine to silver as an auxiliary ligand and increase the abundance of coordination with silver," said Jin-Ping Gao, first author of the study paper and researcher at the Key Laboratory of Magnetic Molecules & Magnetic Information Materials at Shanxi Normal University in Taiyuan, China.

"Here, we introduced raw sodium molybdate... into the reaction process... to guide the synthesis of this quadruple-symmetric 13-silver atom cluster," said Gao. Molybdates are often introduced into the reaction environment and through self-assembly reactions can be obtained as negatively charged molybdenum POMs, which can be encapsulated in a silver cluster structure to change their size, shape and balance charge.

"We isolated an ideal, quadruple-symmetric sandwich cationic (positively charged)... nanocluster, $[\text{Ag}_{13}(\text{MoO}_4)_4(\text{SC}_6\text{H}_4i\text{Pr})_2(\text{dppp})_8]^{3+}$ (Ag_{13}), which is synthesized using four molybdate $[\text{MoO}_4]^{2-}$ anions and protected by two ligands, thiolate ($i\text{PrC}_6\text{H}_4\text{S}^-$) and phosphine (dppp). This work demonstrates that simple molybdate anions also have the potential to induce novel structures, opening the door to the use of this anion to isolate structures of interesting clusters," said Gao.

Nanocluster synthesis is paramount to developing new cluster structures with different chemical and physical properties. "Metal nanoclusters are ideal nanomaterials for applications in different fields, including catalysis, or speeding chemical reactions, and biological applications such as bioimaging, sensing, optics, imaging, biolabeling and targeted drug delivery," said Gao.

The research team looks forward to overcoming new silver [nanocluster](#) synthesis challenges to create the nanomaterials of the future. This work will include experimenting with various protective ligands and understanding how peripheral organic molecules and inorganic cores are bonded to determine their precise chemical structure.

"We aspire to generate more silver clusters with high-symmetry aesthetics using a strategy of mixed-ligand combinations induced by an anionic, or negatively charged, template in the future. Our goal is to enrich the research process of mixing strategies to construct novel silver

clusters and broaden the research progress of molybdate in highly symmetric [silver](#) clusters," said Gao.

More information: Jin-Ping Gao et al, [MoO₄]²⁻-templated D_{4h}-symmetric sandwich Ag₁₃ nanocluster coprotected with thiolate and phosphine, *Polyoxometalates* (2023). [DOI: 10.26599/POM.2023.9140028](#)

Provided by Tsinghua University Press

Citation: Protective molecules facilitate molybdate anion binding to create novel silver nanoclusters (2023, June 20) retrieved 2 May 2024 from <https://phys.org/news/2023-06-molecules-molybdate-anion-silver-nanoclusters.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.