


Previously overlooked algae toxin widespread in southern Indian River Lagoon

June 15 2023



A transmission electron microscopy image of *Pseudo-nitzschia* cells. Credit: Florida Atlantic University

Spanning about one-third of Florida's East coast, the Indian River Lagoon has faced frequent harmful algal blooms in recent years. Among them, *Pseudo-nitzschia* spp., algae that produces the neurotoxin domoic acid.

Domoic acid can bioaccumulate within [food webs](#), causing sickness and death in higher trophic level organisms such as marine mammals and birds, and have been documented in sea turtles in Florida [coastal waters](#) and in bull sharks within the Indian River Lagoon system. In humans, consumption of shellfish contaminated with domoic acid can cause harmful symptoms.

Unlike other harmful algal blooms, Pseudo-nitzschia are not bioluminescent and do not cause water discoloration or large fish kills. Because there are no obvious visual cues of these blooms and monitoring them is event based, documenting their presence is challenging.

Researchers from Florida Atlantic University's Harbor Branch Oceanographic Institute have conducted the first widespread molecular study of domoic acid produced from Pseudo-nitzschia in the Indian River Lagoon. Their results, published in the journal Harmful Algae, suggest domoic acid may be a significant threat to the lagoon's ecosystem.

To gain a better understanding of the ecology of Pseudo-nitzschia in the central and southern portions of the Indian River Lagoon system where monitoring has been less frequent, researchers collected surface water samples from five locations along the lagoon between October 2018 and May 2020. They analyzed 158 samples for phytoplankton concentrations via cell counts that were collected on 68 different dates. Scanning [electron microscopy](#) was conducted on cultured material, which enabled them to morphologically identify six distinct species of Pseudo-nitzschia and compare them with sequencing data.

Results showed Pseudo-nitzschia was present in 87 percent of the samples, which were associated with relatively high salinity waters and cool temperatures. All isolates demonstrated toxicity, and domoic acid was found in 47 percent of surface water samples. Although Pseudo-

nitzschia concentrations were more prominent during the first half of the study, cells were often present at lower concentrations throughout the entire study period. Notable differences between the Indian River Lagoon system and other water bodies where Pseudo-nitzschia blooms occur include water temperatures, water clarity, residence times and salinity.

"The frequent presence of Pseudo-nitzschia, including bloom level concentrations, which we observed in our study, could lead to domoic acid integrating into the food web affecting shellfish, finfish, birds, [sea turtles](#), dolphins, sharks, rays and humans," said Malcolm McFarland, Ph.D., senior author and a research associate at FAU Harbor Branch. "As a nursery for many organisms, and supporting a high amount of biodiversity, the presence of domoic acid could negatively impact the biodiversity of the Indian River Lagoon system."

In addition to a positive relationship with salinity, researchers also found that Pseudo-nitzschia had an inverse relationship with temperature. This suggests Pseudo-nitzschia in the Indian River Lagoon prefers cooler water temperatures for growth. Cell abundances of Pseudo-nitzschia were highest during the late fall, winter and early spring, when temperatures in the lagoon are cooler. Some of the highest abundances of Pseudo-nitzschia in the southern Indian River Lagoon system occurred at sampling sites with the closest proximities to inlets.

"Given the characteristics of the Indian River Lagoon system, such as warm water temperatures, high nutrients, and shallow depths, we think that Pseudo-nitzschia may be a resident population, that is, present year-round," said Stephanie Schreiber, first author and a coordinator of research programs and services at FAU Harbor Branch. "Results from our study suggest Pseudo-nitzschia could be considered a key contributor to phytoplankton community composition in the southern Indian River Lagoon system as well."

As one of multiple toxic [harmful algal blooms](#)-forming organisms within the Indian River Lagoon, understanding its distribution and population dynamics is critical to protecting biodiversity as well as human, animal and ecosystem health.

"Accurate identification methods are critical to determine the presence or absence of toxigenic species of *Pseudo-nitzschia* and to assess the potential threat posed by blooms of these algae," said McFarland.

"Although domoic acid is not present at all times in the southern Indian River Lagoon system, its frequent occurrence suggests it is an important toxin to monitor in the lagoon. Continuous exposures to [domoic acid](#), even at low levels, may have unknown impacts on the [lagoon](#) system over time."

More information: Stephanie Schreiber et al, *Pseudo-nitzschia* species, toxicity, and dynamics in the southern Indian River Lagoon, FL, *Harmful Algae* (2023). [DOI: 10.1016/j.hal.2023.102437](https://doi.org/10.1016/j.hal.2023.102437)

Provided by Florida Atlantic University

Citation: Previously overlooked algae toxin widespread in southern Indian River Lagoon (2023, June 15) retrieved 17 May 2024 from <https://phys.org/news/2023-06-previously-overlooked-algae-toxin-widespread.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--