

How well toxic elements leach out of coal ash depends on the ash's nanoscale composition, shows study

June 6 2023, by Ken Kingery



Leftover sludge from the 2008 coal ash spill at the Kingston TVA power plant. New research indicates that the nanoscale structure of the coal ash plays a large part in whether or not toxic chemicals can leach into the environment from such events. Credit: Helen Hsu-Kim, Duke University

Everyone knows that burning coal causes air pollution that is harmful to the climate and human health. But the ash left over can often be harmful as well.


For example, Duke Energy long stored a liquified form of [coal](#) ash in 36 large ponds across the Carolinas. That all changed in 2014, when a spill at its Dan River site released 27 million gallons of ash pond water into the local environment. The incident raised concerns about the dangers associated with even trace amounts of toxic elements like arsenic and selenium in the ash. Little was known, however, about just how much of these [hazardous materials](#) were present in the ash water or how easily they could contaminate the surrounding environment.

Fears of future spills and seepage caused Duke Energy to agree to pay \$1.1 billion to decommission most of its coal ash ponds over the coming years. Meanwhile, researchers are working on better ways of putting the ash to use, such as recycling it to recover valuable rare earth elements or incorporating it into building materials such as concrete. But to put any potential solution into action, researchers still must know which sources of coal ash pose a hazardous risk due to its chemical makeup—a question that scientists still struggle to answer.

In a new paper published in *Environmental Science: Nano*, researchers at Duke University have discovered that these answers may remain elusive because nobody is thinking small enough. Using one of the newest, most advanced synchrotron light sources in the world—the National Synchrotron Light Source II at Brookhaven National Laboratory—the authors show that, at least for selenium and arsenic, the amount of toxic elements able to escape from coal ash depends largely on their [nanoscale structures](#).

"These results show just how complex coal ash is as a material," said Helen Hsu-Kim, professor of civil and [environmental engineering](#) at Duke University. "For example, we saw arsenic and selenium either attached to the surface of fine grain particles or encapsulated within them, which explains why these elements leach out of some coal ash sources more readily than others."

It's long been known that factors in the surrounding environment such as pH affect how well toxic elements can move from source to surroundings. In previous research, Hsu-Kim showed that the amount of oxygen in a toxin's surroundings can greatly affect its chemistry, and that different sources of coal ash produce vastly different levels of byproducts.



A nanoscale view of two different sources of coal ash shows major differences in how arsenic is part of its makeup. On the left, arsenic atoms (red) coat the surface of a coal ash particle made mainly of iron (green). On the right, the arsenic is encapsulated within the iron particle, making it more difficult for the arsenic to escape. Credit: Helen Hsu-Kim, Duke University

But just because one source of coal ash is high in arsenic doesn't necessarily mean that high amounts of arsenic will leach out of it. Similarly, various sources of ash respond differently to the same environmental conditions. The problem is complex, to say the least. To take a different approach, Hsu-Kim decided to take an even closer look at the source itself.


"Researchers in the field typically use X-ray microscopy with a resolution of one or two micrometers, which is about the same size as the fly ash particles themselves," Hsu-Kim said. "So if a single particle is a single pixel, you're not seeing how the elements are distributed across it."

To shrink these pictures' pixels to the nanoscale, Hsu-Kim turned to Catherine Peters, professor of civil and environmental engineering at Princeton University, and her colleagues to acquire time on the National Synchrotron Light Source II. The futuristic machine creates light beams 10 billion times brighter than the sun to reveal the chemical and atomic structure of materials using light beams ranging from infrared to hard X-rays.

Brookhaven's capabilities were able to provide the researchers a nanoscale map of each particle along with the distribution of elements in each particle. The incredible resolution revealed that coal ash is a compilation of particles of all kinds and sizes.

For example, in one sample the researchers saw individual nanoparticles of selenium that were attached to bigger particles of coal ash, which is a chemical form of selenium that probably isn't very soluble in water. But most of the ash had arsenic and selenium either locked inside individual grains or attached at the surface with relatively weak ionic bonds that are easily broken.

"It was almost like we saw something different in every sample we looked at," Hsu-Kim said. "The wide array of differences really highlights why the main characteristic that we care about—how much of these elements leach out of the ash—varies so much between different samples."



A nanoscale view of two different sources of coal ash shows major differences in how selenium is part of its makeup. On the left, selenium (red) is diffusely distributed around multiple large ash grains made of iron (green) and carbon (blue). On the right, selenium is bound to the surface of ash grains made of iron and carbon, making it more difficult for the selenium to escape. Credit: Helen Hsu-Kim, Duke University

While nobody can say for sure what causes the coal ash to develop its unique composition, Hsu-Kim guesses that it is likely mostly related to how the coal was originally formed millions of years ago. But it might also have something to do with the [power plants](#) that burn the coal. Some plants inject activated carbon or lime into the flue gas, which captures mercury and sulfur emissions, respectively. At 1000 degrees Fahrenheit, toxins such as arsenic and selenium in the flue are gaseous, and the

physics that dictate how the particles will cool and recombine to form ash is uncontrollable.

But regardless of the how, researchers now know that they should be paying closer attention to the fine details encapsulated within the end results.

More information: Nelson A Rivera et al, Nanoscale heterogeneity of arsenic and selenium species in coal fly ash particles: Analysis using enhanced spectroscopic imaging and speciation techniques, *Environmental Science: Nano* (2023). [DOI: 10.1039/D2EN01056A](https://doi.org/10.1039/D2EN01056A)

Provided by Duke University

Citation: How well toxic elements leach out of coal ash depends on the ash's nanoscale composition, shows study (2023, June 6) retrieved 2 May 2024 from <https://phys.org/news/2023-06-toxic-elements-leach-coal-ash.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.