


Asteroid Ryugu's anhydrous ingredients come from afar, study suggests

July 26 2023



A: Ryugu as seen by Hayabusa2. B: The “stone” C0002 collected by Hayabusa2. C: Electron microscope image of a section of C0002. D: Infrared image with anhydrous clasts. E: Spectra of a grain rich in amorphous silicates, of a cometary particle, of the primitive asteroid Hektor and of comet Hale-Bopp. Credit: IAS Orsay

Infrared spectra of anhydrous grains from the carbonaceous asteroid Ryugu indicate a connection between one of the reservoirs from which Ryugu's parent body originated and the reservoirs that formed comets and primitive asteroids in the outer protoplanetary disk.

Ryugu is a second-generation carbonaceous asteroid formed by the reassembly of fragments from a previous larger body in the [main asteroid belt](#). In late 2020, the Japanese mission Hayabusa2 returned to Earth 5.4 g of materials collected on Ryugu. The majority of samples consists of hydrated minerals which formed in Ryugu's parent asteroid by the aqueous alteration of primordial anhydrous grains. These grains reveal the composition of the protoplanetary dust from which Ryugu's parent asteroid formed.

A new study led by IAS (Orsay, France), Tohoku University (Japan), and the SMIS beamline of synchrotron SOLEIL (France) used infrared hyperspectral imaging of two millimeter-sized "stones" of Ryugu to retrieve their mineral composition and to compare Ryugu with other extraterrestrial materials in the lab (meteorites and interplanetary dust) and in space (asteroids and comets). The study is published in *The Astrophysical Journal Letters*.

The infrared analysis of Ryugu's anhydrous grains shows that some of them are rich in amorphous silicates, with a mineral composition similar to that of some anhydrous primitive asteroids, comets, and interplanetary dust particles of cometary origin. These amorphous-rich grains formed by pre-accretional processes in the [protoplanetary disk](#) from which the solar system originated.

Ryugu's parent asteroid would thus be a large planetesimal that formed in the outer solar system from a reservoir close to the accretion region of comets. Planetary migrations would later have transferred Ryugu's parent asteroid to the main asteroid belt. Aqueous alteration would then

have determined the spectral diversity of "primitive" asteroid classes we observe today.

More information: R. Brunetto et al, Ryugu's Anhydrous Ingredients and Their Spectral Link to Primitive Dust from the Outer Solar System, *The Astrophysical Journal Letters* (2023). [DOI: 10.3847/2041-8213/acdf5c](https://doi.org/10.3847/2041-8213/acdf5c)

Provided by IAS Orsay

Citation: Asteroid Ryugu's anhydrous ingredients come from afar, study suggests (2023, July 26) retrieved 2 May 2024 from

<https://phys.org/news/2023-07-asteroid-ryugu-anhydrous-ingredients-afar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.