


Researchers study gut-to-CNS translocation of silver nanomaterials

July 13 2023, by Liu Jia



Working model of the transneuronal transport of NPs from the gut to the CNS. When Ag NPs enter the gastrointestinal tract by oral gavage, they adsorb proteins to form a nano-corona that facilitates the uptake of NPs by various cell subsets of the gut, especially enterocytes and enteric nerve cells in submucosal/myenteric plexus. Thereafter, the vagus nerves act as direct conduits to mediate the transneuronal transport of Ag NPs along the gut-brain axis. Meanwhile, the spinal nerves mediate the transneuronal transport of Ag NPs along the gut-spinal

cord axis. Credit: *Science Advances* (2023). DOI: 10.1126/sciadv.adg2252

Recently, a research team led by Prof. Chen Chunying from the National Center for Nanoscience and Technology (NCNST) of the Chinese Academy of Sciences (CAS) revealed that peripheral nerve fibers act as direct conduits for silver nanomaterials translocation from the gut to the central nervous systems (CNS). The study was published in *Science Advances*.

The [transport](#) route determines the final target organ and subsequent biological effects of exogenous nanoparticles (NPs). The specific transfer route map of NPs is crucial to unveil their nanotoxicological as well as nanomedical potential. However, there are still substantial knowledge gaps in understanding of the transport of NPs from the gut to the CNS, which hinders both the understanding of the gut-CNS axis communication and the development of new therapeutic strategies that target CNS-associated diseases.

In this study, the researchers first revealed transfer routes for Ag NMs along the gut-CNS axis in addition to [blood circulation](#). They demonstrated both theoretically and experimentally the direct translocation of Ag NMs from the gut to the CNS (both the brain and [spinal cord](#)) via peripheral [nerve](#) fibers. Using [laser ablation](#) and field flow fractionation inductively coupled plasma mass spectrometry, they showed that Ag NMs administered by oral gavage became enriched in a particle state in the brain and spinal cord of mice while not efficiently entering the blood.

In addition, the researchers found that the vagus and spinal nerves mediate the transneuronal transport of Ag NMs along the gut-brain axis and gut-spinal cord axis, respectively, with specific gut cell subsets

(enterocytes and enteric nerve cells) taking up significant levels of Ag NMs for transfer to the connected peripheral nerves using Cy-TOF analysis.

The results implicated that transneuronal transport as a mechanism for the direct trafficking of NPs, filling a crucial gap in understanding the behavior of NPs in biological systems.

This study integrated a variety of methods to break through the "classical pathway" of blood circulation mediated NPs transport. For the first time, it proposed that peripheral nerve fibers as the "non-classical pathway" mediate NPs transport between organs in the body. This path of NPs in the living body provides important theoretical support for the development of nanomedicines targeting CNS-related diseases.

More information: Xiaoyu Wang et al, Peripheral nerves directly mediate the transneuronal translocation of silver nanomaterials from the gut to central nervous system, *Science Advances* (2023). [DOI: 10.1126/sciadv.adg2252](https://doi.org/10.1126/sciadv.adg2252)

Provided by Chinese Academy of Sciences

Citation: Researchers study gut-to-CNS translocation of silver nanomaterials (2023, July 13) retrieved 29 April 2024 from <https://phys.org/news/2023-07-gut-to-cns-translocation-silver-nanomaterials.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.