

Researchers propose mechanism for heattransfer enhancement in convective turbulence

July 24 2023



(a) The Nusselt number ratio Nu/Nu_{RB} and (b) the Reynolds number ratio Re/Re_{RB} versus the wave number kH, respectively. In (a) and (b), different



symbols represent different combinations of (Ra, *A/H*) as indicated in the legend of (a). (c) The viscous and thermal boundary-layer thicknesses versus the Rayleigh number for the RB flow. Credit: *Physical Review Letters* (2023). DOI: 10.1103/PhysRevLett.130.204001

Researchers from the Department of Mechanics and Engineering Science at Peking University (PKU), led by Prof. Chen Shiyi and Prof. Yang Yantao, have recently proposed a mechanism to enhance the efficiency of heat transfer in convective turbulence. The paper has been published in *Physical Review Letters*, entitled "Boundary-Layer Disruption and Heat-Transfer Enhancement in Convection Turbulence by Oscillating Deformations of Boundary."

Convective turbulence, driven by gravity-induced instabilities resulting from temperature gradients, is a prevalent occurrence in various natural flows, including the atmosphere, ocean, and mantle. It is referred to as an indispensable process for both heat and mass transport and finds practical applications in <u>heat dissipation</u> in devices and indoor temperature control.

However, the existing methods for regulating convective turbulence suffer from several drawbacks. They only function in a limited range of parameters, demand an extremely high frequency, and consequently, the overall heat transfer enhancement is restricted.

The PKU research team's novel approach focuses on introducing oscillating deformations in the form of standing waves in a flow system where the top is at a high temperature and the bottom at a low temperature, thus effectively modulating transport efficiency.

The research results demonstrate that significant increases in heat flux



can be achieved only when the amplitude of boundary oscillation approaches the thickness of the boundary layers. Notably, the dimensionless <u>heat flux</u> more than doubles when the dimensionless frequency precisely reaches 8. In essence, oscillating deformations of boundary can break the bottleneck of classic thermal convection in the <u>boundary layer</u> and open a new venue for modulating the convection turbulence.

More information: Leiqi Yuan et al, Boundary-Layer Disruption and Heat-Transfer Enhancement in Convection Turbulence by Oscillating Deformations of Boundary, *Physical Review Letters* (2023). <u>DOI:</u> <u>10.1103/PhysRevLett.130.204001</u>

Provided by Peking University

Citation: Researchers propose mechanism for heat-transfer enhancement in convective turbulence (2023, July 24) retrieved 10 May 2024 from <u>https://phys.org/news/2023-07-mechanism-heat-transfer-convective-turbulence.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.