

Fire in northern Australia's tropical savanna is a threat to endangered fairy-wrens

July 31 2023, by Niki Teunissen and Anne Peters



Credit: Niki Teunissen/AWC, Author provided

Wildfire threatens the survival of endangered [purple-crowned fairy-wrens](#) living along the rivers and creeks of northern Australia, our new research has found.

For [almost two decades](#), we studied the fairy-wrens at a [wildlife sanctuary](#) in the far north of Western Australia.

Over this time, one low-intensity [fire](#) and one high-intensity fire burnt through our study site. Both occurred late in the [wet season](#), when fires generally burn at lower intensity. But drought and [weather conditions](#) meant the second fire unexpectedly burnt at high intensity instead.

We wanted to find out what happened to the birds before, during and after each fire. We found even low-intensity burns reduced population density. As this species is a [biological indicator](#) of ecosystem health, our [new research](#) can help fine-tune fire management practices, to reduce the extent and intensity of fires along waterways.

Fire in the tropics

Fire is particularly [common](#) in tropical monsoonal savanna. The vegetation thrives during the wet season, then dries out over the dry season. This creates plenty of fuel late in the dry season, leading to frequent fires.

Deliberately introducing fire in the early dry season, when fires generally burn at low intensity, can reduce large intense wildfires later in the year. So fire management is often used for [conservation](#) and [carbon farming](#).

But approaches to fire management that best protect "riparian" communities are relatively poorly understood. Riparian zones are the strips of vegetation along creeks and rivers. They play an important role in tropical savanna landscapes. They support a highly [diverse range of species](#), provide corridors for animals to move through the landscape, and form a [cool refuge](#) from heat and drought.

Unfortunately they are also particularly [sensitive to fire](#), making it ever more urgent to better protect these key places.

Understanding how riparian fire affects the species that depend on

waterway vegetation for their entire life cycle is a good place to start.

Fairy-wrens and fire

We study a population of 200–300 [purple-crowned fairy-wrens](#) (*Malurus coronatus coronatus*) along 15km of waterways.

Each bird in this population has been tagged with a unique small colored leg band. This enables us to recognize individuals and follow them throughout their life.

We gather detailed information on bird survival, movement and reproduction. This is key to quantifying how—and to what extent—fire impacts populations.

In our study we found both low-intensity and high-intensity fire reduced the number of fairy-wrens in the burnt areas for at least two and a half years. The effect of high-intensity fire was much stronger, reducing the number of fairy-wrens by half.

Next, we investigated what mechanism caused these declines. We showed birds did not move out of burnt habitat, probably because they live in such well-defined territories year-round.

Instead, we found the low-intensity fire reduced breeding success by 80% during and shortly after the fire.

The [high-intensity](#) fire caused a decline in wrens through a different mechanism. We found birds in the fire-affected area were no more likely to die during the fire itself than birds in adjacent unburnt areas. Yet, they were 30% more likely to die over the next two to eight months after the fire.

This is probably because the quality of the riparian habitat was greatly reduced by this fire, which may have made it harder for the birds to find food, cover from predators, or find protection from the heat in subsequent months.

Protecting riparian zones from fire

Wildfires are becoming more [frequent and severe](#) as [climate change](#) worsens. These changes are [transforming natural systems](#) and threatening the diversity of life on Earth. We saw this in Australia in 2019–20, when the Black Summer fires pushed many species closer to extinction.

More frequent and severe fire is forecast for riparian zones, for various reasons. For example, extended droughts as well as large flood events (which deposit woody debris as fuel for fire), [increase the risk](#) of severe riparian fires. Additionally, riparian strips can become a corridor for fire under certain conditions.

The challenge for land managers is to minimize the impact of fire when it eventually enters the riparian zone. We suggest fire management can be used to reduce the extent and intensity of riparian fires. In particular, we recommend introducing low-intensity burns parallel and perpendicular to riparian zones so they have minimal impact yet create breaks along these riparian corridors, to prevent large sections burning at once.

Our study indicates the high sensitivity of [riparian zones](#) to fire, even when fire occurs during the wet season and burns at low intensity. Our findings call for more consideration by fire managers of the effects of fire on riparian habitat, and for further research to enhance our understanding of savanna riparian fire biology.

This article is republished from [The Conversation](#) under a Creative

Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Fire in northern Australia's tropical savanna is a threat to endangered fairy-wrens (2023, July 31) retrieved 30 April 2024 from <https://phys.org/news/2023-07-northern-australia-tropical-savanna-threat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.