

Using new tech to identify the cause of an old phenomenon in meat tenderness

July 10 2023, by Anna Zarra Aldrich



Credit: Fermo Sergio, Pixabay

When you bite into a juicy steak, tough isn't what you're looking for. There's a lot of science behind the spectrum from tough to tender.



Chaoyu Zhai, assistant professor of animal science in the College of Agriculture, Health and Natural Resources, recently published a paper in the *Journal of Agricultural and Food Chemistry* using machine learning algorithms to explain how the activity of an enzyme responsible for meat tenderness is modified on the molecular level.

This is the first study to do so in the field of meat science.

An enzyme called calpain, specifically calpain-1, is known as the "meat tenderization enzyme." Calpain-1 cuts other proteins in the meat and makes these <u>protein</u> fragments more water-soluble. Having more fragmented protein molecules in meat makes it more tender.

While scientists have known about calpain-1's role in meat tenderness since the 1990s, they did not know much about how it can be modified.

"Protein modification is like a person wearing different clothes every day," Zhai says. "It's still the same 'person' but depending on the outfit, it may be better prepared to tackle different jobs."

Scientists noticed that when meat is packaged with a higher concentration of oxygen, which helps preserve color, the meat was tougher, but they didn't know why.

Zhai and his collaborators discovered that the presence of compounds known as <u>lipid</u> peroxidation products can modify calpain-1 activity, explaining this mystery.

"We provided a new explanation for something we see a lot," Zhai says. "By studying this, we help other researchers understand similar events."

Lipid peroxidation occurs when lipids—such as cell membranes and fatty acids—are exposed to oxygen. Free radicals "steal" electrons from



lipids in cell membranes, causing cell damage. For example, lipid peroxidation occurs as perishable food gets older, causing it to degrade and smell.

In this paper, Zhai focused on three lipid peroxidation products: malondialdehyde (MDA), hexenal, and 4-hydroxynonenal (HNE), which have been previously identified in meat.

These compounds have three, six, and nine carbons, respectively. The more carbon molecules a compound has, the more hydrophobic, or water repelling, they tend to be. This means that these compounds likely inhibit calpain activity at different levels.

"When the carbon chain number increases, those compounds tend to be more hydrophobic," Zhai says. "They will potentially cause more functional changes to the protein."

Zhai worked with Steven M. Lonergan and Elisabeth J. Huff-Lonergan, two experts on calpains from Iowa State University. The research team used artificial intelligence algorithms to understand structural changes corresponding to functional changes in the protein.

Normally, to determine the 3D structure of a protein, scientists will freeze it and quickly capture an X-Ray image. Machine learning algorithms allow researchers to simply input a <u>protein sequence</u> and determine its structure, a much quicker and less expensive process.

"By using these algorithms, we're able to understand our results more effectively and allow us to better interpret it," Zhai says.

This approach can be used in any field interested in protein activity—a long and diverse list of disciplines.



Zhai found that HNE, the most hydrophobic compound his group tested, severely decreased calpain-1 activity but, unexpectedly, did not totally eliminate it. Another compound called hexenal had a similar effect, but the reduction in activity was less dramatic. Some compounds like MDA, surprisingly, increased calpain-1 activity at the lowest concentration the team tested.

"That's definitely surprising," Zhai says. "We would normally just assume if it affects activity, we would see a decrease rather than an increase."

The team also discovered that calpain-1 had already begun undergoing modification in freshly cut muscle that had not been exposed to the air.

"People never looked for it," Zhai says. "They assume because this type of modification involves oxygen, it happens later, when the meat is exposed to oxygen. Turns out it happens way earlier than people think."

In another field, researchers found that lipid peroxidation is related to natural aging, while high calpain-1 activity is related to many human diseases, like heart failure and Alzheimer's disease.

The body's metabolism is imperfect and generates byproducts, like <u>lipid</u> <u>peroxidation</u> products. These byproducts bind to proteins and modify them. As they accumulate over a lifetime, the effects of aging are more and more visible.

"This direction of the research is definitely worth pursuing because it's more common than we thought," Zhai says. "These were high-quality commercial pigs, they are raised to be healthy and strong, and they still withstood a lot of <u>protein modification</u> caused by lipid oxidation products, so it's very interesting."



The next step of this research is to study what other enzymes are modified by compounds like MDA and HNE in <u>meat</u> production.

"With more results, you have more research questions," Zhai says. "If you don't look for them, you don't find them."

More information: Chaoyu Zhai et al, Lipid Peroxidation Products Influence Calpain-1 Functionality In Vitro by Covalent Binding, *Journal of Agricultural and Food Chemistry* (2023). DOI: 10.1021/acs.jafc.3c01225

Provided by University of Connecticut

Citation: Using new tech to identify the cause of an old phenomenon in meat tenderness (2023, July 10) retrieved 10 May 2024 from <u>https://phys.org/news/2023-07-tech-phenomenon-meat-tenderness.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.