

Researchers seek to determine climate drivers of early human interbreeding

August 10 2023



Photo of the remaining Denisova 11 (Denny) bone fragment from Denisova Cave in Russia, that comes from a daughter to a Neanderthal mother and a Denisovan father. Credit: Katerina Douka, Tom Higham, Institute for Basic



Science

A new study published in the journal *Science* by an international team finds that past changes in atmospheric CO_2 and corresponding shifts in climate and vegetation played a key role in determining when and where early human species interbred.

Modern-day people carry in their cells a small quantity of DNA deriving from other human species, namely the Neanderthals and the elusive Denisovans.

Back in 2018, scientists announced to the world the discovery of an individual, later nicknamed Denny, who lived 90,000 years ago and who was identified as a daughter to a Denisovan father and a Neanderthal mother (Slon et al. 2018). Denny, along with fellow mixed-ancestry individuals found at Denisova cave, testifies that interbreeding was probably common among hominins, and not limited to Homo sapiens.

To unravel when and where human hybridization took place, scientists usually rely on paleo-genomic analysis of extremely rare fossil specimens and their even scarcer ancient DNA content.

In the new *Science* paper, the team of climate experts and paleobiologists from South Korea and Italy pursued a different approach. Using existing paleo-anthropological evidence, genetic data and supercomputer simulations of past climate, the team found that Neanderthals and Denisovans had different environmental preferences. More specifically, Denisovans were much more adapted to cold environments, characterized by <u>boreal forests</u> and even tundra, compared to their Neanderthal cousins who preferred <u>temperate forests</u> and grassland.



"This means that their habitats of choice were separated geographically, with Neanderthals typically preferring southwestern Eurasia and Denisovans the northeast," says Dr. Jiaoyang Ruan, postdoctoral researcher at the IBS Center for Climate Physics (ICCP), South Korea and lead author of the study.

However, according to their realistic computer simulations the scientists found that in warm interglacial periods, when Earth's orbit around the sun was more elliptic and northern hemisphere summer occurred closer to the sun, the hominin habitats began to overlap geographically. "When Neanderthals and Denisovans shared a common habitat, there were more encounters and interactions among the groups, which would have increased the chance of interbreeding," adds Prof. Axel Timmermann, corresponding author of the study and director of the ICCP and professor at Pusan National University.





Illustration of Neanderthal (redscale) /Denisovan (greenscale) preferred habitats. Potential interbreeding areas in Central Asia and northern Europe are indicated by overlapping colors and baby-shapes. Credit: Institute for Basic Science

The simulation of past habitat overlaps does not only put the first generation Neanderthal/Denisovan hybrid Denny into a climatic context, but it also agrees with other known episodes of interbreeding ~78, 120 thousand years ago. Future paleo-genetic reconstructions can be used to test the robustness of the new supercomputer model-based predictions of potential interbreeding intervals around 210 and 320 thousand years ago.



To further determine the climate drivers of the east-west interbreeding seesaw, the scientists looked more closely at how vegetation patterns changed over Eurasia during the past 400 thousand years.

They discovered that elevated atmospheric CO_2 concentrations and mild interglacial conditions caused an eastward expansion of temperate forest into central Eurasia which created dispersal corridors for Neanderthals into Denisovan lands. "It is as if glacial-interglacial shifts in climate created the stage for a unique and long-lasting human love story, whose genetic traces are still visible today," comments Dr. Ruan.

One of the key challenges the researchers faced in their study was to estimate the preferred climatic conditions for Denisovans. "To deal with the very sparse Denisovan dataset, we had to devise new statistical tools, which could also account for known ancestral relationships among <u>human species</u>," says Prof. Pasquale Raia from University of Naples, Federico II in Italy, co-author of the study.





Multiple models demonstrate how glacial cycle variability synchronized hominin habitat overlap, Denisovan-Neanderthal interbreeding, and early human gene flow. Credit: Axel Timmermann

"This allowed us for the first time to estimate where Denisovans could have lived. To our surprise, we found that, apart from areas in Russia and China, also northern Europe would have been a suitable environment for them," he adds.

Whether Denisovans ever lived west of the Altai mountains is unknown; but it can be tested using large-sample genetic analyses of Denisovan ancestry in European populations. Such analysis is expected to shed new light on the relationship between early dispersal, habitat encroachment and human genetic diversification.

More information: Jiaoyang Ruan et al, Climate shifts orchestrated hominin interbreeding events across Eurasia, *Science* (2023). DOI: <u>10.1126/science.add4459</u>. <u>www.science.org/doi/10.1126/science.add4459</u>

Provided by Institute for Basic Science

Citation: Researchers seek to determine climate drivers of early human interbreeding (2023, August 10) retrieved 15 May 2024 from <u>https://phys.org/news/2023-08-climate-drivers-early-human-interbreeding.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.