

Research suggests daily temperature difference is main factor affecting rubber latex yield in Xishuangbanna

August 3 2023, by Zhang Nannan







Credit: Pixabay/CC0 Public Domain

Rubber trees (Hevea brasiliensis) are a major source of rubber latex. Xishuangbanna is the largest rubber growing region in China. However, Xishuangbanna's climate is considered sub-optimal for rubber cultivation. Rubber latex yield is influenced by several factors, including temperature, disease, other biotic conditions, and plantation management. Quantifying and clarifying of the relationships between leaf flushing phenology, powdery mildew disease, temperature, and latex yield will inform future strategies to improve latex yield.

In a study published in *International Journal of Biometeorology*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) of the Chinese Academy of Sciences (CAS) and the Kunming Institute of Botany of CAS examined the effects of temperature, phenology, and powdery mildew disease on rubber latex yield in March using <u>observational data</u> on daily rubber latex yield combined with detailed phenology, powdery mildew, and temperature data from 2004 to 2010 at a state-owned farm in the Xishuangbanna, Yunnan, China.

The researchers found that the daily temperature difference from January to March was the most important factor influencing rubber latex yield, surpassing both rubber phenology and powdery mildew disease. In addition, rubber phenology during defoliation and re-foliation (November to March) was found to be critical for rubber latex yield.

Daily temperature variation had both direct and indirect effects on rubber latex yield. Temperature indirectly influenced yield by altering the duration of leaf development to maturation and exposure it to



powdery mildew disease.

"This study is the most comprehensive quantitative assessment of the relationship and the potential linkages between rubber latex yield and rubber leaf defoliation phenology, <u>powdery mildew</u> disease, and <u>temperature</u>," said Zhai Deli of XTBG.

The researchers recommended that future experiments on <u>rubber</u> latex yield, latex physiology and leaf diseases should cover both defoliation and re-foliation periods.

More information: De-Li Zhai et al, Rubber latex yield is affected by interactions between antecedent temperature, rubber phenology, and powdery mildew disease, *International Journal of Biometeorology* (2023). DOI: 10.1007/s00484-023-02515-2

Provided by Chinese Academy of Sciences

Citation: Research suggests daily temperature difference is main factor affecting rubber latex yield in Xishuangbanna (2023, August 3) retrieved 13 May 2024 from <u>https://phys.org/news/2023-08-daily-temperature-difference-main-factor.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.