

Researchers attach electrodes to individual atomically precise graphene nanoribbons

August 15 2023, by Anna Ettlin



Empa researchers and their international collaborators have successfully attached carbon nanotube electrodes to individual atomically precise nanoribbons. Credit: Empa

Graphene nanoribbons have outstanding properties that can be precisely controlled. Researchers from Empa and ETH Zurich, in collaboration with partners from Peking University, the University of Warwick and the Max Planck Institute for Polymer Research, have succeeded in



attaching electrodes to individual atomically precise nanoribbons, paving the way for precise characterization of the fascinating ribbons and their possible use in quantum technology.

Quantum technology is promising, but also perplexing. In the coming decades, it is expected to provide us with various technological breakthroughs: smaller and more precise sensors, highly secure communication networks, and powerful computers that can help develop new drugs and materials, control financial markets, and predict the weather much faster than current computing technology ever could.

To achieve this, we need so-called quantum materials: substances that exhibit pronounced quantum <u>physical effects</u>. One such material is <u>graphene</u>. This two-dimensional structural form of carbon has unusual physical properties, such as extraordinarily high tensile strength, thermal and electrical conductivity—as well as certain <u>quantum effects</u>. Restricting the already two-dimensional material even further, for instance, by giving it a ribbon-like shape, gives rise to a range of controllable quantum effects.

This is precisely what Mickael Perrin's team leverage in their work. For several years now, scientists in Empa's Transport at Nanoscale Interfaces laboratory, headed by Michel Calame, have been conducting research on graphene nanoribbons under Perrin's leadership. "Graphene nanoribbons are even more fascinating than graphene itself," explains Perrin. "By varying their length and width, as well as the shape of their edges, and by adding other atoms to them, you can give them all kinds of electrical, magnetic, and optical properties."



OI MALIGIUN DOID



Anticu et al toto OVIO



The extremely narrow ribbons with their atomically precise edge exhibit strong quantum effects, making them particularly interesting to researchers. Credit: Empa

Ultimate precision—down to single atoms

Research on the promising ribbons isn't easy. The narrower the ribbon, the more pronounced its quantum properties are—but it also becomes more difficult to access a single ribbon at a time. This is precisely what must be done in order to understand the unique characteristics and possible applications of this quantum material and distinguish them from collective effects.

In a new study published in the journal Nature Electronics, Perrin and



Empa researcher Jian Zhang, together with an international team, have succeeded for the first time in contacting individual long and atomically precise graphene nanoribbons. "A graphene nanoribbon that is just nine <u>carbon atoms</u> wide measures as little as 1 nanometer in width," Zhang says. To ensure that only a single nanoribbon is contacted, the researchers employed electrodes of a similar size. They used carbon nanotubes that were also only 1 nanometer in diameter.

Precision is key for such a delicate experiment. It begins with the source materials. The researchers obtained the graphene nanoribbons via a strong and long-standing collaboration with Empa's nanotech@surfaces laboratory, headed by Roman Fasel. "Roman Fasel and his team have been working on graphene nanoribbons for a long time and can synthesize many different types with atomic precision from individual precursor molecules," Perrin explains. The precursor molecules came from the Max Planck Institute for Polymer Research in Mainz.

As is often required for advancing the state of the art, interdisciplinarity is key, and different international research groups were involved, each bringing in their own specialty to the table. The carbon nanotubes were grown by a research group at Peking University, and to interpret the results of the study, the Empa researchers collaborated with computational scientists at the University of Warwick. "A project like this would not be possible without collaboration," Zhang emphasizes.

Contacting individual ribbons by nanotubes posed a considerable challenge for the researchers. "The carbon nanotubes and the graphene nanoribbons are grown on separate substrates," Zhang explains. "First, the nanotubes need to be transferred to the device substrate and contacted by metal electrodes. Then we cut them with high-resolution electron-beam lithography to separate them into two electrodes." Finally, the ribbons are transferred onto the same substrate. Precision is key: Even the slightest rotation of the substrates can significantly reduce the



probability of successful contact. "Having access to high-quality infrastructure at the Binnig and Roher Nanotechnology Center at IBM Research in Rüschlikon was essential to test and implement this technology," Perrin says.



The properties of nanoribbons vary depending on their width and the shape of their edges. Credit: Empa

From computers to energy converters

The scientists confirmed the success of their experiment through charge transport measurements. "Because quantum effects are usually more



pronounced at low temperature, we performed the measurements at temperatures close to absolute zero in a high vacuum," Perrin explains. He adds, "Due to the extremely small size of these nanoribbons, we expect their quantum effects to be so robust that they are observable even at room temperature."

This, the researcher says, could allow us to design and operate chips that actively harness quantum effects without the need for an elaborate cooling infrastructure.

"This project enables the realization of single nanoribbon devices, not only to study fundamental quantum effects such as how electrons and phonons behave at the nanoscale, but also to exploit such effects for applications in quantum switching, quantum sensing, and quantum energy conversion," adds Hatef Sadeghi, a professor at the University of Warwick who collaborated on the project.

Graphene nanoribbons are not ready for commercial applications just yet, and there is still a lot of research to be done. In a follow-up study, Zhang and Perrin aim to manipulate different quantum states on a single <u>nanoribbon</u>. In addition, they plan on creating devices based on two ribbons connected in series, forming a so-called double quantum dot.

Such a circuit could serve as a qubit—the smallest unit of information in a quantum computer. Perrin plans to explore the use of nanoribbons as highly-efficient energy converters.

More information: Jian Zhang et al, Contacting individual graphene nanoribbons using carbon nanotube electrodes, *Nature Electronics* (2023). DOI: 10.1038/s41928-023-00991-3



Provided by Swiss Federal Laboratories for Materials Science and Technology

Citation: Researchers attach electrodes to individual atomically precise graphene nanoribbons (2023, August 15) retrieved 12 May 2024 from <u>https://phys.org/news/2023-08-electrodes-individual-atomically-precise-graphene.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.