


Hawaii's climate future: Increasingly dry regions increase fire risk while wet areas get wetter

August 16 2023, by Matthew McCartney, Kevin Hamilton



The top map shows the long-term average rainfall rate observed in Hawaii. The bottom map displays the result of the computer simulation, showing how closely the rain shadows are reproduced. Credit: [Adapted from Zhang et al., Journal of Climate, American Meteorological Society](#)

The islands of Hawaii are world renowned for their generally pleasant and tranquil weather. However, the Aug. 8, 2023, wildfire tragedy on Maui was a stark reminder that Hawaii also can experience drought and hot, dry, windy weather, providing the conditions for destructive fires.

Hawaii has seen a generally [rising trend](#) in the amount of [land that burns](#) each year as the [local climate warms](#). Climate change was [one of several contributors](#) to Maui's wildfire catastrophe, and rising temperatures and associated [rainfall](#) changes are expected to increase the islands' fire risk. These changing weather patterns will also affect Hawaii's ecosystems and freshwater resources.

[I am a meteorologist](#) at the University of Hawaii, and I have worked with colleagues to develop sophisticated [computer climate simulations](#) that [project local rainfall changes](#) over the 21st century.

Our results suggest that as the planet warms, Hawaii's dry regions will get drier, heightening the fire risk. At the same time, its wet areas will become wetter.

The [drier parts of the state](#) in particular have reason to be concerned about the [future of fresh water](#) available for residential, commercial and agricultural uses. In addition, changes in rainfall are expected to affect the [distribution of plants in Hawaii](#), harming some unique native species [such as silversword](#) and increasing some [invasive grass species](#) that enhance fire danger.

Average rainfall drops sharply in the rain shadow


While Hawaii is home to some of the wettest spots on earth, it also has regions that receive little [rain](#).

The very steep mountains on each of the main Hawaiian islands block

the prevailing northeast trade winds. This results in abundant rain on the slopes facing the windward direction and dry "rain shadows" in the leeward areas. Maui's west coast tourist communities, including Lahaina, are in one of those rain shadows.

Hawaii is remarkable for its exceptionally strong gradations in the [average rainfall](#) rates over very short distances. The summit of Mt. Waialeale in central Kauai, known as one of the rainiest places on Earth, [receives an average of about 450 inches of rain per year](#). The town of Kehaka, 15 miles to the southwest and in the rain shadow, [receives less than 20 inches per year](#) on average.

These sharp differences over short distances have made projecting future [climate change](#) in Hawaii a particularly daunting challenge.



The change in average rainfall rates projected to occur over the 21st century as simulated in the computer model. These changes are expressed as a percentage of present-day rainfall rates at each location. Credit: Adapted from Zhang et al., *Journal of Climate*, American Meteorological Society

Computer [models used to project the future climate](#) approximate the atmospheric wind, temperature and humidity at discrete points on a regular grid. The horizontal spacing between grid points in [global climate models](#) is [typically 20 miles or larger](#). To put that in perspective, Maui is

only [26 miles by 48 miles](#) at its widest.

Predicting Hawaii's changing climate

We created a model that [zooms in on Hawaii](#) and is able to capture those variations, including the rain shadow effect.

Using that model, we [simulated the Hawaiian climate](#) at the end of the 21st century under a scenario in which [global greenhouse gas emissions](#) from human activities continue at a rate that drives a global increase of temperature of about 4 degrees Fahrenheit (2.2 C). Such a scenario is quite plausible and even demands some significant reduction in current emission rates, but still pushes well beyond the internationally agreed [goal of keeping global warming under 3.6 F \(2 C\)](#) compared with preindustrial levels.

We found that in the wet windward areas of Hawaii, rainfall is projected to increase substantially. That includes increasingly frequent extreme downpours. On the other hand, rainfall is predicted to decrease substantially over much of the rain shadow regions.

This overall "wet gets wetter and dry gets drier" trend [is generally found in global model projections of climate warming](#). Our computer model shows that it also applies to the rainfall gradations over the very short distances relevant for Hawaii.

The "dry gets drier" aspect is particularly important for formulating Hawaii's plans to adapt to climate change. The soil in already dry regions may become even drier as rainfall decreases and warmer air promotes more evaporation from the surface. That includes Maui's highly developed west coast and agricultural areas that until recently [were home to large sugar cane farms](#).

Developing better projections to help prepare

Rainfall rates will still vary year to year. Indeed, Hawaii rainfall is known to display [quite strong year-to-year and even decade-to-decade variations](#), due in large part to the influences of [El Niño](#) and the [Pacific decadal oscillation](#), both natural climate patterns. Such natural variations are expected to coexist with the overall century-long trend toward drier or wetter conditions.

Simulating rainfall in climate models still has many uncertainties, and there are particular challenges in representing the fine geographical details of the rain in Hawaii. Another [study using a different approach](#) produced results broadly consistent with ours but projects an overall stronger drying trend in the islands.

While further research should help reduce the uncertainties in [climate](#) projections, our results suggest that, in the long term, Hawaii needs to prepare for more extreme conditions, including a heightened risk of wildfires.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Hawaii's climate future: Increasingly dry regions increase fire risk while wet areas get wetter (2023, August 16) retrieved 10 May 2024 from <https://phys.org/news/2023-08-hawaii-climate-future-dry-regions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.