

The modern sea spider had started to diversify by the Jurassic, study finds

August 17 2023



Palaeopycnogonides gracilis (normal colour). Credit: Dr Romain Sabroux

An extremely rare collection of 160-million-year-old sea spider fossils from Southern France are closely related to living species, unlike older fossils of their kind.

These fossils are very important to understand the evolution of sea spiders. They show that the diversity of sea spiders that still exist today had already started to form by the Jurassic.

Lead author Dr. Romain Sabroux from the University of Bristol's School of Earth Sciences, said, "Sea spiders (Pycnogonida), are a group of marine animals that is overall very poorly studied. "New insights into the sea [spider](#) fauna (Arthropoda: Pycnogonida) of La Voulte-sur-Rhône, France (Jurassic: Callovian)" was published in *Papers in Palaeontology*.


"However, they are very interesting to understand the evolution of arthropods [the group that includes insects, arachnids, crustaceans, centipedes and millipedes] as they appeared relatively early in the arthropod tree of life. That's why we are interested in their evolution.

"Sea spider fossils are very rare, but we know a few of them from different periods. One of the most remarkable fauna, by its diversity and its abundance, is the one of La Voulte-sur-Rhône that dates back to the Jurassic, some 160 million years ago."

Unlike older sea spider fossils, the La Voulte pycnogonids are morphologically similar (but not identical) to living species, and previous studies suggested they could be closely related to living sea spider families. But these hypotheses were restricted by the limitation of their observation means. As it was impossible to access what was hidden in

the rock fossils, Dr. Sabroux and his team traveled to Paris and set out to investigate this question with cutting-edge approaches.

Dr. Sabroux explained, "We used two methods to reinvestigate the morphology of the fossils: X-ray microtomography, to 'look inside' the rock, find morphological features hidden inside and reconstruct a 3D model of the fossilized specimen; and reflectance transformation imaging, a picture technic that relies on varied orientation of the light around the [fossil](#) to enhance the visibility of inconspicuous features on their surface.



Palaeopycnogonides gracilis in blue and pink ; that is drawn from the reflectance transformation Imaging technic). Credit: Dr Romain Sabroux

"From these new insights, we drew new morphological information to compare them with extant species," explained Dr. Sabroux.

This confirmed that these fossils are close relatives to surviving pycnogonids. Two of these fossils belong to two living pycnogonid families: *Colossopantopodus boissinensis* was a Colossendeidae while another, *Palaeoendeis elmii* was an Endeidae. The third species, *Palaeopycnogonides gracilis*, seems to belong to a family that has disappeared today.


"Today, by calculating the difference between the DNA sequences of a sample of species, and using DNA evolution models, we are able to estimate the timing of the evolution that bind these species together," added Dr. Sabroux.

"This is what we call a molecular clock analysis. But quite like a real clock, it needs to be calibrated. Basically, we need to tell the clock: 'we know that at that time, that group was already there.' Thanks to our work, we now know that Colossendeidae, and Endeidae were already 'there' by the Jurassic."

Now, the team can use these minimal ages as calibrations for the molecular clock, and investigate the timing of Pycnogonida evolution. This can help them understand, for example, how their diversity was impacted by the different biodiversity crises that distributes over the Earth history.

They also plan to investigate other pycnogonid fossil faunae such as the fauna of Hunsrück Slate, in Germany, which dates from the Devonian, some 400 million years ago.

With the same approach, they will aim to redescribe these species and understand their affinities with extant species; and finally, to replace in the tree of life of Pycnogonida all the pycnogonid fossils from all periods.



Palaeopycnogonides gracilis - the 3D model drawn from the CT-scan data.
Credit: Dr Romain Sabroux

Dr. Sabroux added, "These fossils give us an insight of sea spiders living 160 million years ago."

"This is very exciting when you have been working on the living pycnogonids for years.

"It is fascinating how these pycnogonids look both very familiar, and very exotic. Familiar, because you can definitely recognize some of the families that still exist today, and exotic because of small differences like the size of the legs, the length of the body, and some other [morphological characteristics](#) that you do not find in modern [species](#)."

More information: New insights into the sea spider fauna (Arthropoda: Pycnogonida) of La Voulte-sur-Rhône, France (Jurassic: Callovian), *Papers in Palaeontology* (2023).

Provided by University of Bristol

Citation: The modern sea spider had started to diversify by the Jurassic, study finds (2023, August 17) retrieved 8 May 2024 from <https://phys.org/news/2023-08-modern-sea-spider-diversify-jurassic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--