


Newly discovered fungus helps destroy a harmful food toxin

August 31 2023



Patulin toxicity is a serious food safety hazard, and scientists are looking for ways to control it. Now, researchers from TUS, Japan, have identified, for the first time, a mold strain that can convert patulin into desoxypatulinic acid and other less toxic substances, with potential applications for patulin biocontrol. Credit: *MicrobiologyOpen* (2023). DOI: 10.1002/mbo3.1373

Patulin ($C_7H_6O_4$), a mycotoxin produced by several types of fungi, is toxic to a variety of life forms, including humans, mammals, plants, and microorganisms. In particular, environments lacking proper hygienic measures during food production are susceptible to patulin contamination as many of these fungi species tend to grow on damaged or decaying fruits, specifically apples, and even contaminate apple products, such as apple sauce, apple juice, jams, and ciders.

Responsible for a wide variety of health hazards, including nausea, lung congestion, ulcers, intestinal hemorrhages, and even more serious outcomes, such as DNA damage, immunosuppression, and increased cancer risk, patulin toxicity is a serious concern worldwide. As a result, many countries have imposed restrictions on the permitted levels of patulin in [food products](#), especially baby foods as infants are more vulnerable to the effects of patulin.

Treatment of patulin toxicity include oxygen therapy, immunotherapy, detoxification therapy, and nutrient therapy. However, as prevention is often better than cure, scientists have been on the lookout for efficient ways to mitigate patulin toxicity in food products.

To this end, a research team including Associate Professor Toshiki Furuya from Tokyo University of Science (TUS) in Japan, recently screened for soil microorganisms that can potentially help keep patulin toxicity in check. Their study, published in *MicrobiologyOpen*, was co-authored by Ms. Megumi Mita, Ms. Rina Sato, and Ms. Miho Kakinuma, all from TUS.

The team cultured microorganisms from 510 [soil samples](#) in a patulin-rich environment, looking for those that would thrive in presence of the toxin. Next, in a second screening experiment, they used high-performance liquid chromatography (HPLC) to determine the survivors that were most effective in degrading patulin into other less harmful

chemical substances. Accordingly, they identified a filamentous fungal (mold) strain, *Acremonium* sp. or "TUS-MM1," belonging to the genera *Acremonium*, that fit the bill.

The team then performed various experiments to shed light on the mechanisms by which TUS-MM1 degraded patulin. This involved incubating the mold strain in a patulin-rich solution and focusing on the substances that gradually appeared both inside and outside its cells in response to patulin over time.

One important finding was that TUS-MM1 cells transformed any absorbed patulin into desoxypatulinic acid, a compound much less toxic than patulin, by adding hydrogen atoms to it. "When we started this research, only one other filamentous fungal strain had been reported to degrade patulin," comments Dr. Furuya. "However, prior to the present study, no degradation products had ever been identified. In this regard, to our knowledge, TUS-MM1 is the first filamentous fungus shown to be capable of degrading patulin into desoxypatulinic acid."

Moreover, the team found that some of the compounds secreted by TUS-MM1 cells can also transform patulin into other molecules. By mixing patulin with the extracellular secretions of TUS-MM1 cells and using HPLC, they observed various degradation products generated from patulin.

Encouragingly, experiments on *E. coli* bacterium cells revealed that these products are significantly less toxic than patulin itself. Through further chemical analyses, the team showed that the main agent responsible for patulin transformation outside the cells was a thermally stable but highly reactive compound with a low molecular weight.

Overall, the findings of this study take us a step closer toward efficient solutions for controlling the levels of patulin in food. Dr. Furuya says,

"Elucidating the pathways via which microorganisms can degrade patulin would be helpful not only for increasing our understanding of the underlying mechanisms in nature but also for facilitating the application of these organisms in biocontrol efforts."

More information: Isolation and characterization of filamentous fungi capable of degrading the mycotoxin patulin, *MicrobiologyOpen* (2023).
[DOI: 10.1002/mbo3.1373](https://doi.org/10.1002/mbo3.1373)

Provided by Tokyo University of Science

Citation: Newly discovered fungus helps destroy a harmful food toxin (2023, August 31)
retrieved 21 May 2024 from <https://phys.org/news/2023-08-newly-fungus-destroy-food-toxin.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--