


# Oil eating microbes reshape droplets to optimize biodegradation

August 18 2023, by Bob Yirka



Bright-field sequences showing biofilm phenotypes. (A) Time-lapse sequence showing the formation of a monolayer on a trapped drop. The edge of the drop trap is indicated by the white dashed line. The confluent monolayer is formed at  $t_0$  ( $t = 0$  h). (B) Development of a spherical biofilm (SB) on an oil drop. The oil drop radius monotonically decreases in time. At 12 h, the droplet is outlined with a cyan line and the biofilm is outlined by a magenta line, as a guide for the reader. (C) Time-lapse sequence showing the development of a dendritic biofilm (DB) on an oil drop. The biofilm deforms the surface, initially generating wrinkles and protrusions, that fragmenting the droplet into tube-like segments and numerous smaller irregularly shaped volumes of oil covered with cells at later times. Credit: *Science* (2023). DOI: 10.1126/science.adf3345

A team of French and Japanese environmental scientists has found that one kind of oil-eating microbe reshapes droplets to optimize biodegradation. In their study, reported in the journal *Science*, the group isolated *Alcanivorax borkumensis* bacteria specimens in a lab setting, fed them crude oil, and then watched how they worked together to eat the oil as quickly and efficiently as possible. Terry McGenity and Pierre Philippe Laissue with the University of Essex's School of Life Sciences have published a Perspective piece in the same journal issue describing the work.

Prior research has shown that there are many microbes living in the ocean that feed on oil, eventually cleaning away oil not cleaned up by human efforts. Prior research has also shown that such microbes are not able to consume crude oil until it disperses into droplets, which can take a long time. In this new effort, the researchers sought to learn more about the process of crude oil consumption by sea microbes. To that end, they collected *A. borkumensis* specimens and tested them in their lab.

Under a [microscope](#), the research team observed that *A. borkumensis* formed biofilms around oil droplets—but they did so in two major ways. In one experiment, *A. borkumensis* samples that had not been exposed to crude oil before were introduced to simple crude oil [droplets](#). Groups of the bacteria converged on a droplet, forming a [sphere](#). The sphere shape persisted until the entire oil droplet had been consumed.

But when the team exposed samples with experience consuming [crude oil](#), their behavior was much more advanced. Initially, upon converging on a droplet, a sphere formed—but then finger-like protrusions formed, radiating out from the sphere, each completely covered with bacteria. The result was much faster, more efficient consumption of the droplet.

The researchers suggest that the formation of the protrusions results in more oil surface area exposure, allowing more of the [bacteria](#) to consume the oil droplet at the same time, compared to the simple sphere, resulting in faster consumption.

**More information:** M. Prasad et al, *Alcanivorax borkumensis* biofilms enhance oil degradation by interfacial tubulation, *Science* (2023). [DOI: 10.1126/science.adf3345](https://doi.org/10.1126/science.adf3345).  
[www.science.org/doi/10.1126/science.adf3345](https://www.science.org/doi/10.1126/science.adf3345)

Terry J. McGenity et al, Bacteria stretch and bend oil to feed their appetite, *Science* (2023). [DOI: 10.1126/science.adj4430](https://doi.org/10.1126/science.adj4430)

© 2023 Science X Network

Citation: Oil eating microbes reshape droplets to optimize biodegradation (2023, August 18) retrieved 11 May 2024 from <https://phys.org/news/2023-08-oil-microbes-reshape-droplets-optimize.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.