

## Plankton evidence reveals a seasonally icefree ocean during the Last Interglacial

August 4 2023



Photograph taken in the marginal ice zone of the Arctic Ocean from the Swedish icebreaker Oden, summer 2021. Credit: Flor Vermassen

A subpolar species associated with Atlantic water expanded far into the Arctic Ocean during the Last Interglacial, analysis of microfossil content



of sediment cores reveals. This implies that summers in the Arctic were ice free during this period. The findings are published in *Nature Geoscience*.

Arctic sea ice, an important component of the Earth system, is disappearing fast under climate warming. Summer sea ice is anticipated to vanish entirely within this century. To gain a deeper understanding of the climate dynamics in a world without Arctic sea ice, researchers have turned to analogs from the geological past.

"The Last Interglacial, between 129,000 and 115,000 years BP, is an interesting <u>period</u> to study because it is the last time in Earth's history when global average temperatures were similar or perhaps higher than currently and sea levels were considerably higher, up to +6 to +9 m," said Flor Vermassen, postdoctoral researcher at Stockholm University.

However, the extent of sea ice during this period has been intensely debated and there is no consensus, limiting understanding of this period and the ability of researchers to simulate it in <u>climate models</u>.

To address this, a team of marine geology researchers from the Department of Marine Geological Sciences at Stockholm University analyzed the microfossil content of an array of sediment cores from sites that today lie directly beneath the thickest parts of the modern Arctic ice pack.

In these cores, they investigated the variability in the occurrence and composition of planktonic foraminifera, a type of free-floating, shell-building unicellular zooplankton that is sensitive to changes in oceanographic and <u>environmental conditions</u>.





Scanning electron microscope images of the planktonic foraminifer Turborotalita quinqueloba. Scale bars = 100 micron. Credit: Stockholm University

The researchers found high abundances of the typically subpolar, Atlantic-water species Turborotalita quinqueloba, documenting a largescale expansion of the species far into the central Arctic Ocean. The ecological preference of T. quinqueloba for predominantly ice-free, seasonally productive waters typically present in the Atlantic Ocean suggests that it was following similar conditions that had spread to the central Arctic Ocean.

The absence of summer sea ice and increased influence of Atlantic currents in the Arctic domain during the Last Interglacial are analogous to <u>ocean</u> transformations being observed today in parts of the Arctic, and collectively referred to as "Atlantification" of the Arctic Ocean.

"The finding that the Arctic Ocean was seasonally ice-free during the Last Interglacial is worrying because this period would have been only around 1.5°C above pre-industrial levels, comparable to the targets of the Paris Agreement. Yet the global sea level is estimated to have been several meters higher than at present," said Flor Vermassen.



Hence, the researchers propose the Last Interglacial as the most recent and potentially most relevant geological epoch for investigating a seasonally ice-free Arctic Ocean, particularly if the objectives of the Paris Agreement are not exceeded.

"To fully comprehend the <u>physical conditions</u> and environment of this unfamiliar Arctic during the Last Interglacial, additional quantitative proxy reconstructions of sea-surface temperature and other water mass parameters are needed, along with targeted climate and oceanographic model studies of the same period," said Flor Vermassen.

**More information:** Flor Vermassen et al, A seasonally ice-free Arctic Ocean during the Last Interglacial, *Nature Geoscience* (2023). DOI: 10.1038/s41561-023-01227-x

## Provided by Stockholm University

Citation: Plankton evidence reveals a seasonally ice-free ocean during the Last Interglacial (2023, August 4) retrieved 11 May 2024 from <u>https://phys.org/news/2023-08-plankton-evidence-reveals-seasonally-ice-free.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.