

Tree mortality in the Black Forest on the rise, climate change is key driver: Study

August 23 2023



Forest dieback in the Black Forest. Credit: Heinrich Spiecker

Climate impacts such as dry, hot summers reduce the growth and increase the mortality of trees in the Black Forest because they negatively influence the climatic water balance, i.e., the difference between precipitation and potential evapotranspiration. That is the central finding of a long-term study of the influence of climate and climate change on trees in the Black Forest conducted by Prof. Dr. Hans-Peter Kahle and Prof. Dr. Heinrich Spiecker, who are both professors of forest growth and dendroecology at the University of Freiburg.

The researchers used a consistent time series of 68 years (1953 to 2020) as the basis for their research. It covers the annual mortality of all trees in an almost 250-thousand-hectare area of the public forests in the Black Forest. They analyzed these data in comparison with a second data series on the climatic water balance for the months of May to September. The results of their research have been published in the scientific journal *Global Change Biology*.


Tree mortality peaked in 2019

"Our time series on [tree growth](#) and mortality in the Black Forest is unique and allows quantitative analysis of the impacts of heat and drought," says Kahle. Together with his colleague, he examined [trees](#) that died as a result of insect or fungal infestation, atmospheric deposition, frost or drought, and of other causes.

The time series on climatic water balance—which covers 140 years (1881–2020)—shows a continuous downward trend. "Also reflected in this analysis is the [tree mortality](#) at the end of the last century known as Forest Dieback I. In the Black Forest, its extent can be explained solely

by [weather conditions](#) and the associated bark beetle infestation," explains Spiecker.

At that time, however, a maximum of 12% of the sustainable annual growth died off, yet the mortality rate after extremely dry summers in recent years has increased to 40% of sustainable annual growth. Sustainable growth describes the sum of the average annual growth of wood which is used as a basis for calculating the annual allowable sustainable timber harvest.



Time series of observed and modeled tree mortality over time in the Black Forest. The observed mortality from 1953 to 2020 is presented in red bars. The blue bars describe the modeled mortality from 1881 to 2020 based on the climatic water balance, and the green bars describe the modeled mortality from 1900 to 2020 based on the annual radial growth rate index. Credit: Heinrich Spiecker

In 2019, mortality reached a peak at more than seven times the average

mortality rate in the period from 1953 to 2017. Kahle elaborates, "What is also striking is a certain regularity in the occurrence of cool-humid and warm-dry periods that repeated themselves in the past around every fourteen years. But we've observed that the cool-humid periods are becoming weaker while the warm-dry periods are getting more severe."

More information: Heinrich Spiecker et al, Climate-driven tree growth and mortality in the Black Forest, Germany—Long-term observations, *Global Change Biology* (2023). [DOI: 10.1111/gcb.16897](https://doi.org/10.1111/gcb.16897)

Provided by University of Freiburg

Citation: Tree mortality in the Black Forest on the rise, climate change is key driver: Study (2023, August 23) retrieved 29 April 2024 from <https://phys.org/news/2023-08-tree-mortality-black-forest-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.