


Atomic-scale spin-optical laser: New horizon for optoelectronic devices

September 6 2023



The spin-valley optical microcavity is constructed by interfacing an inversion-

asymmetric (yellow core region) and an inversion-symmetric (cyan cladding region) photonic spin lattice. By virtue of a photonic Rashba-type spin splitting of a bound state in the continuum, this heterostructure enables a selective lateral confinement of the emergent photonic spin-valley states inside the core for high-Q resonances. Consequently, the coherent and controllable spin-polarized lasing (red and blue beams) is achieved from valley excitons in an incorporated WS₂ monolayer (purple region). Credit: Scholar designer co, LTD

Researchers at the Technion—Israel Institute of Technology have developed a coherent and controllable spin-optical laser based on a single atomic layer. This discovery is enabled by coherent spin-dependent interactions between a single atomic layer and a laterally confined photonic spin lattice, the latter of which supports high-Q spin-valley states through the photonic Rashba-type spin splitting of a bound state in the continuum.

Published in *Nature Materials* and also featured in the journal's Research Briefing, the achievement paves the way to study coherent spin-dependent phenomena in both classical and quantum regimes, opening new horizons in [fundamental research](#) and optoelectronic devices exploiting both electron and photon spins.

Can we lift the spin degeneracy of light sources in the absence of magnetic fields at room temperature? According to Dr. Rong, "Spin-optical light sources combine photonic modes and electronic transitions and therefore provide a way to study the exchange of spin information between electrons and photons and to develop advanced optoelectronic devices."

"To construct these sources, a prerequisite is to lift the spin degeneracy between the two opposite spin states either in their photonic or electronic parts. This is usually accomplished by applying magnetic fields under a

Faraday or Zeeman effect, although these approaches generally require strong magnetic fields and cannot produce miniaturized sources. Another promising way takes advantage of artificial magnetic fields for photonic spin-split states in momentum space, underpinned by a geometric phase mechanism."

"Unfortunately, previous observations of spin-split states have relied heavily on propagation modes with low quality factors, which impose undesired limitations on spatial and temporal coherence of the sources. This approach is also hindered by the spin-controllable properties of a bulk laser gain material being unavailable or nontrivial to access for active control of the sources, especially in the absence of magnetic fields at room temperature."

To achieve high-Q spin-split states, the researchers constructed photonic spin lattices with different symmetry properties, which comprise an inversion-asymmetry core and inversion-symmetry cladding integrated with a WS₂ monolayer to create laterally confined spin-valley states. The essential inversion-asymmetry lattice the researchers use has two important properties.

1. A controllable spin-dependent reciprocal lattice vector due to space-variant geometric phases from its constituting inhomogeneous-anisotropic nanoholes. This vector splits a spin-degenerate band into two spin-polarized branches in momentum space, being referred to as the photonic Rashba effect.
2. A pair of high-Q symmetry-enabled (quasi-) bound states in the [continuum](#), that is, $\pm K$ (corners of the Brillouin zone) photonic spin-valley states, at the band edges of the spin-split branches. Moreover, the two states form a coherent superposition state with equal amplitudes.

Professor Koren noted that, "We used a WS₂ monolayer as the gain

material because this direct-bandgap transition metal dichalcogenide possesses unique valley pseudospins, which have been widely investigated as an alternative information carrier in valleytronics. Specifically, their $\pm K'$ valley excitons (radiated as in-plane spin-polarized dipole emitters) can be selectively excited by spin-polarized light according to a valley-contrasted selection rule, thus enabling active control of spin-optical light sources without magnetic fields."

In the monolayer-integrated spin-valley microcavities, $\pm K'$ valley excitons couple to $\pm K$ spin-valley states owing to polarization matching, and spin-optical excitonic lasing is achieved at room temperatures through strong optical feedback. Meanwhile, $\pm K'$ valley excitons (initially without a phase correlation) are driven by the lasing mechanism to find the minimum-loss state of the system, which leads them to re-establish a phase-locked correlation according to the opposite geometric phases of $\pm K$ spin-valley states.

This lasing-mechanism-driven valley coherence removes the need for cryogenic temperatures to suppress the intervalley scattering. Moreover, the minimum-loss state of the Rashba monolayer laser can be regulated to be satisfied (broken) via a linear (circular) pump polarization, which provides a way to control the lasing intensity and spatial coherence.

"The unveiled photonic spin valley Rashba effect provides a general mechanism to construct surface-emitting spin-optical light sources. The demonstrated valley coherence in the monolayer-integrated spin-valley microcavity makes a step towards achieving entanglement between $\pm K'$ valley excitons for quantum information by means of qubits," explains Professor Hasman.

"For a long time, our group has been working on developing spin optics to harness photonic spin as an effective tool to control the behavior of electromagnetic waves. In 2018, we were attracted by valley pseudospins

in two-dimensional materials, and therefore began a long-term project to study the active control of atomic-scale spin-optical light sources in the absence of magnetic fields. We initially tackled the challenge of coherent geometric phase pickup from individual valley excitons by using a non-local Berry-phase defect mode."

"However, the underlying coherent addition of multiple valley excitons of the realized Rashba monolayer light sources remained unsolved, owing to the lack of a strong synchronizing mechanism between the excitons. This issue inspired us to think about high-Q photonic Rashba modes. Following innovations in new physical approaches, we achieved the Rashba monolayer laser described here."

The achievement paves the way to study coherent spin-dependent phenomena in both classical and quantum regimes, opening new horizons in fundamental research and optoelectronic devices exploiting both electron and photon spins.

The study was conducted in the research group of Professor Erez Hasman, head of the Atomic-Scale Photonics Laboratory, in collaboration with Professor Elad Koren, head of the Laboratory for Nanoscale Electronic Materials and Devices in the Department of Materials Science and Engineering, and Professor Ariel Ismach at Tel Aviv University.

The two groups at the Technion are in association with the Helen Diller Quantum Center and Russell Berrie Nanotechnology Institute (RBNI). Dr. Kexiu Rong conducted and led the research, and collaborated with Dr. Xiaoyang Duan, Dr. Bo Wang, Dr. Vladimir Kleiner, Dr. Assael Cohen, Dr. Pranab K. Mohapatra, Dr. Avinash Patsha, Dr. Subhrajit Mukherjee, Dror Reichenberg, Chieh-li Liu, and Vladi Gorovoy.

The fabrication was performed at the Micro-Nano Fabrication &

Printing Unit (MNF&PU) of the Technion.

More information: Kexiu Rong et al, Spin-valley Rashba monolayer laser, *Nature Materials* (2023). [DOI: 10.1038/s41563-023-01603-3](https://doi.org/10.1038/s41563-023-01603-3)

A spin-optical monolayer laser based on a photonic spin lattice, *Nature Materials* (2023). [DOI: 10.1038/s41563-023-01623-z](https://doi.org/10.1038/s41563-023-01623-z)

Provided by Technion - Israel Institute of Technology

Citation: Atomic-scale spin-optical laser: New horizon for optoelectronic devices (2023, September 6) retrieved 2 May 2024 from <https://phys.org/news/2023-09-atomic-scale-spin-optical-laser-horizon-optoelectronic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.