

Researchers develop novel catalyst for carbon dioxide hydrogenation to formate

September 12 2023, by Li Yuan



Edge-rich MoS_2 is presented as a promising catalyst for CO_2 hydrogenation to formate with superior activity and stability. The selective formation of formate



is enabled by using surface OH* and H* species from H_2O dissociation on the edge-sulfur vacancies as moderate hydrogenating agents. Credit: *Angewandte Chemie International Edition* (2023). DOI: 10.1002/anie.202307086

Hydrogenation of carbon dioxide (CO_2) to formate is an attractive approach for the utilization of this greenhouse gas. However, nonprecious metal-based catalysts for CO_2 hydrogenation to formate suffer from either low activity or low stability. It is still challenging to develop low-cost and high-performance catalysts.

Recently, a research group led by Prof. Deng Dehui from the Dalian Institute of Chemical Physics (DICP) of the Chinese Academy of Sciences (CAS) has developed an edge-rich molybdenum disulfide (ER- MoS_2) catalyst for CO₂ hydrogenation to formate with superior activity and high <u>stability</u>.

The study was published in <u>Angewandte Chemie International Edition</u> on July 20.

In the study, the ER-MoS₂ with abundant edges delivered a high turnover frequency of 780.7 h^{-1} with formate selectivity of over 99% at 200°C, and exhibited good stability.

The researchers revealed that sulfur vacancies at MoS_2 edges were the <u>active sites</u>, and the selective production of formate was enabled via a new water-mediated hydrogenation mechanism, in which surface OH* and H* species from H₂O dissociation on the edge-sulfur vacancies served as moderate hydrogenating agents with residual O* reduced by H₂.

"This work opens new avenues for developing low-cost non-noble metal



catalysts for the hydrogenation of CO_2 to formate," said Prof. Deng. "The water-mediated reaction mechanism also provides insights for designing MoS_2 -based catalysts for selective hydrogenation reactions."

More information: Zifeng Wang et al, Boosting CO2 Hydrogenation to Formate over Edge-Sulfur Vacancies of Molybdenum Disulfide, *Angewandte Chemie International Edition* (2023). DOI: 10.1002/anie.202307086

Provided by Chinese Academy of Sciences

Citation: Researchers develop novel catalyst for carbon dioxide hydrogenation to formate (2023, September 12) retrieved 21 May 2024 from <u>https://phys.org/news/2023-09-catalyst-carbon-dioxide-hydrogenation-formate.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.