


New cosmological constraints on the nature of dark matter

September 7 2023



Dark matter fluctuations in the lens system MG J0414+0534. The whitish blue color represents the gravitationally lensed images observed by ALMA. The calculated distribution of dark matter is shown in orange; brighter regions indicate higher concentrations of dark matter and dark orange regions indicate lower concentrations. Credit: ALMA(ESO/NAOJ/NRAO), K. T. Inoue et al.


New research, published in *The Astrophysical Journal*, has revealed the distribution of dark matter in never-before-seen detail, down to a scale of 30,000 light-years. The observed distribution fluctuations provide better constraints on the nature of dark matter.

Mysterious dark matter accounts for most of the matter in the universe. Dark matter is invisible and makes itself know only through its [gravitational effects](#). Dark matter has never been isolated in a laboratory, so researchers must rely on "natural experiments" to study it.

One type of natural experiment is a [gravitational lens](#). Sometimes by random chance, two objects at different distances in the universe will lie along the same line-of-sight when seen from Earth. When this happens, the spatial curvature caused by the matter around the foreground object acts like a lens, bending the path of light from the background object and making a lensed image. However, it is difficult to achieve the high resolution to detect clumps of dark matter that are less massive than [galaxies](#) in natural experiments, so the exact nature of dark matter has been poorly constrained.

A team of Japanese researchers led by Professor Kaiki Taro Inoue at Kindai University used ALMA (Atacama Large Millimeter/submillimeter Array) to study the gravitational lens system known as MG J0414+0534 in the direction of the constellation Taurus. In this system, the foreground object forms not one, but four images of

the background object, due to the gravitational force of a massive galaxy acting on the light.



A conceptual diagram of the gravitational lens system MG J0414+0534. Dark matter associated with the lensing galaxy is shown in pale blue and white. Dark matter in intergalactic space is shown in orange. Solid lines show the actual paths of the radio waves which are bent by gravity. Dotted lines show the apparent observed positions of the lensed images. Credit: NAOJ, K. T. Inoue

With the help of the bending effect and their new data analysis method, the team was able to detect fluctuations in the dark matter distribution along the line-of-sight in higher resolution than ever before, down to a scale of 30,000 light-years.

The new constraints provided by the observed distribution are consistent

with models for slow-moving or "cold" dark matter particles.

In the future, the team plans to further constrain the nature of [dark matter](#) with additional observations.

More information: ALMA Measurement of 10 kpc-scale Lensing Power Spectra towards the Lensed Quasar MG J0414+0534, *The Astrophysical Journal* (2023). [DOI: 10.3847/1538-4357/aceb5f](https://doi.org/10.3847/1538-4357/aceb5f)

Provided by National Institutes of Natural Sciences

Citation: New cosmological constraints on the nature of dark matter (2023, September 7)
retrieved 2 May 2024 from
<https://phys.org/news/2023-09-cosmological-constraints-nature-dark.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.