

Helicopter-based observations uncover warm ocean flows toward Totten Ice Shelf in Southeast Antarctica

September 11 2023



The Totten Ice Shelf. Credit: Yoshihiro Nakayama

An international team of scientists has successfully conducted large-scale helicopter-based observations along the coast of East Antarctica and has identified pathways through which warm ocean water flows from the open ocean into ice shelf cavities for the first time.

During six days of observations, the team was able to retrieve [temperature](#) and other data at 67 sites covering the entire continental shelf region off the Totten Ice Shelf, the floating portion of the Totten Glacier. The study enabled the team to obtain an overall picture of warm [ocean](#) water flowing into the ice shelf cavity. The paper is published in the journal *Geophysical Research Letters*.


The Totten Glacier has attracted global attention because of the sheer scope of the environmental impacts if the glacier were to melt entirely: global sea levels could be raised by nearly four meters. It is thus necessary to observe the entire continental shelf region to understand the pathways and mechanisms of warm water mass inflow. However, such observations have been historically hindered by the region's intense sea ice and icebergs.

The team, including Assistant Professor Yoshihiro Nakayama and Associate Professor Shigeru Aoki of Hokkaido University's Institute of Low Temperature Science, conducted helicopter-based observations with cooperation by Japan's Maritime Self-Defense Force as part of the 61st Japanese Antarctic Research Expedition (JARE61).

Researchers took a CH-101 helicopter from the icebreaker Shirase to observation sites where they dropped two kinds of ocean observation sensors: AXCTD (Airborne eXpendable Conductivity, Temperature, and Depth) and AXBT (Airborne Bathy-Thermograph). The team was able to investigate underwater temperatures and salinity using data from the sensors.

The team found that broad ocean heat—stretching across the 150-kilometer range between the longitudes of 116.5° east and 120.5° east—was intruding toward the shelf. Seawater with a temperature of about 1°C higher than the freezing temperature was observed between 50 and 100 meters above the seabed. This showed, for the first time, the

overall picture of warm water mass flowing toward the Totten Ice Shelf.



A satellite photo of the region surrounding the Totten Glacier, showing the Totten Ice Shelf and Moscow University Ice Shelf, with sampling paths (orange lines), sampling locations (yellow and red circles), and the 1,000 m depth contour (cyan outlines). Credit: *Geophysical Research Letters* (2023). DOI: 10.1029/2022GL097864

This study also demonstrated the efficacy of using helicopters in conducting such research. They are cheaper and faster than operating an icebreaker, have superb maneuverability, and are not affected by sea ice conditions. The identified pathways can be observed in detail by icebreakers.

More information: Yoshihiro Nakayama et al, Helicopter-Based

Ocean Observations Capture Broad Ocean Heat Intrusions Toward the Totten Ice Shelf, *Geophysical Research Letters* (2023). [DOI: 10.1029/2022GL097864](https://doi.org/10.1029/2022GL097864)

Provided by Hokkaido University

Citation: Helicopter-based observations uncover warm ocean flows toward Totten Ice Shelf in Southeast Antarctica (2023, September 11) retrieved 21 May 2024 from <https://phys.org/news/2023-09-helicopter-based-uncover-ocean-totten-ice.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--