

# Most nitrogen from farms goes into the 'nitrogen blanket' and direct deposition is very local: Report

#### September 12 2023



Atmospheric NH<sub>3</sub> concentration (left panel) and bulk deposition (right panel), plotted for each sampling moment. Sampling moments that had farming practices are highlighted by the vertical lines, the practices can be seen at the labels on the x-axes (AF = Artificial Fertilizer; AF Manure = Application of manure). Temperature is plotted in red (left plot, right y-axis), and precipitation is plotted in blue (right plot, right y-axis). In the boxplots, mean values are indicated by the crosses, and median values are indicated by the lines. Credit: *Nitrogen deposition around dairy farms: Spatial and temporal patterns* (2023).

#### On behalf of the Mesdag Zuivelfonds, researchers from the University



of Amsterdam have measured within which radius around dairy farms the nitrogen emitted by the farm is deposited. By far the largest part of the emitted nitrogen ( $\approx 90\%$ ) ends up in higher air layers of the atmosphere and will precipitate elsewhere. Within the measured area, with a radius of 500 meters around the farm, most of the remainder ( $\approx$ 10%) precipitates within a radius of 100 meters.

<u>These results</u> are in accordance with previous research and the models from the Dutch RIVM (National Institute for Public Health and the Environment).

The University of Amsterdam (UvA) was asked to conduct this research in response to questions from farmers' organizations regarding the spatial distribution of various <u>nitrogen</u> sources (agriculture, traffic, aviation) in the nitrogen deposition models used by the Dutch authorities. The research focused on two <u>dairy farms</u>. One <u>farm</u> was located in a more urban area, with a lot of traffic in the area; the other was more isolated and in a rural area.

### **Highest deposition on farmyard**

Several sub-studies have been carried out under the leadership of Albert Tietema of the UvA's Institute for Biodiversity and Ecosystem Dynamics (IBED). He and his colleagues used both existing methods to measure nitrogen deposition and a couple of new, experimental methods, including the use of bio-indicators. Together, these methods provided a clear picture.

Only 9% of the nitrogen emitted by a farm ends up within a radius of 500 meters from the stable. At that distance, the farm's contribution has fallen to a quarter of the total nitrogen concentration in the air and to a fifth of the nitrogen deposition. Within these 500 meters, nitrogen deposition decreases exponentially as the distance from the farm



increases. Nitrogen that precipitates in the vicinity of the farm is mainly deposited on the farmyard itself, within a radius of 100 meters around the farm.

### Nitrogen blanket

Nitrogen is also deposited outside 500 meter radius, but at this distance, an individual farm can no longer be identified as the most important source. Nitrogen that precipitates here mainly comes from the so-called "nitrogen blanket." That term refers to the total nitrogen in the atmosphere that covers our country like a blanket, and that comes from a multitude of both agricultural and industrial sources.

### Model and measurements

According to the researchers, their results are in line with other studies. The team also compared how well their findings correspond with the predicted spatial distribution of nitrogen deposition around the farms based on the RIVM's OPS model. The model appeared to slightly underestimate how much nitrogen ends up within 15 meters of a stable, but beyond that the predictions appeared to agree well with the direct measurements.

Tietema says, "We saw that the predictions became more accurate as more characteristics of the specific farm were included in the model, instead of based on a standard model farm. For example, the number and size of the openings through which nitrogen emissions occur is an important factor to take into consideration."

## Methods

The following methods were used in the various sub-studies:



- Biweekly measurement of ammonia concentration in the air with filters fitted with an acid that captures ammonia
- A funnel/bottle system to determine precipitation of nitrogen such as ammonium and ammonia (wet deposition and part dry deposition)
- Sampling of wet <u>deposition</u> only, consisting a funnel/bottle fitted with a lid that opens when it rains
- New method: automated flux chambers that almost continuously measure ammonia exchange between plants and the atmosphere, for example also after fertilization
- New methods: bio-monitors such as special ryegrass, locally present mosses and algae present in ditches. Dry mass increase is a measure of <u>nitrogen deposition</u> and the ratio of isotopes of nitrogen is an indication of the type of nitrogen source.

**More information:** Report: <u>www.uva.nl/binaries/content/as ...</u> <u>emporal-patterns.pdf</u>

#### Provided by University of Amsterdam

Citation: Most nitrogen from farms goes into the 'nitrogen blanket' and direct deposition is very local: Report (2023, September 12) retrieved 17 May 2024 from <u>https://phys.org/news/2023-09-nitrogen-farms-blanket-deposition-local.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.