


New pathway of diatom-mediated calcification and its impact on the biological pump

September 25 2023



It was discovered that the photosynthesis of *S. costatum* can induce substantial aragonite precipitation from artificial/natural seawater under significantly lower supersaturation levels required for the precipitation of inorganic CaCO_3 . Credit: Science China Press

A research team found that the photosynthesis of *Skeletonema costatum* (*S. costatum*), a common diatom species, can induce substantial aragonite precipitation from artificial/natural seawater under significantly lower supersaturation levels required for the precipitation

of inorganic CaCO_3 .

Researchers have discovered that during the growth process of *S. costatum*, there is a significant decrease in total alkalinity (TA) and $[\text{Ca}^{2+}]$ in the bulk medium. The precipitated white particles were confirmed to be aragonite crystals through X-ray diffraction. Scanning [electron microscope images](#) revealed that the diatom cells were enveloped by spherical crystals with diameters ranging from 40 to 70 μm , forming aggregates of *S. costatum* and aragonite.

Further investigations found that this extracellular calcification process is primarily driven by the combined effect of elevated extracellular CO_3^{2-} concentration and the adsorption and aggregation of Ca^{2+} during photosynthesis. This enables *S. costatum* to induce substantial aragonite precipitation at significantly lower supersaturation levels than those required for inorganic CaCO_3 precipitation.

The team also observed TA deviation from the conservative mixing during *S. costatum* blooms in the East China Sea. This further supports the possibility of a new diatom-mediated calcification pathway occurring in the ocean.

This breakthrough finding has [profound implications](#) for our understanding of oceanic [carbon](#) cycling. Diatoms are the most important primary producers and organic carbon transporters in the ocean. The newly discovered diatom-mediated extracellular calcification pathway may establish a novel connection between the particle inorganic carbon pump and the organic carbon pump.

On one hand, the release of CO_2 during the extracellular calcification process is considered as "counter carbonate pump." However, in the diatom-mediated extracellular calcification process, due to the maintenance of high pH in the water, the released CO_2 may be more

readily absorbed by algae, rather than being released into the atmosphere.

On the other hand, the [calcification](#), through the formation of aggregates of diatoms and aragonite, enhances the efficiency of organic carbon sinking and increases the transport capacity of the biological carbon pump.

This study not only changes our understanding of carbon cycling in [marine ecosystems](#) but also provides new perspectives for the ocean carbon cycle research.

The findings are published in the journal *Science Bulletin*. This study was led by associate professor Yiwen Pan (Institute of Ocean College, Zhejiang University).

More information: Yiwen Pan et al, New pathway of diatom-mediated calcification and its impact on the biological pump, *Science Bulletin* (2023). [DOI: 10.1016/j.scib.2023.08.020](https://doi.org/10.1016/j.scib.2023.08.020)

Provided by Science China Press

Citation: New pathway of diatom-mediated calcification and its impact on the biological pump (2023, September 25) retrieved 17 May 2024 from <https://phys.org/news/2023-09-pathway-diatom-mediated-calcification-impact-biological.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--