

# How does the social behavior of wheat plants influence grain production?

September 6 2023

---



Researchers from IPK Leibniz Institute have investigated how the behavior of an individual wheat plant under limiting light conditions influences the performance of the whole community. Credit: IPK Leibniz Institute/ T. Schnurbusch

One of the most significant drivers of crop evolution stems from the changes in the selection associated with the shift of plants from a highly heterogeneous and biodiverse natural environment into a homogeneous monoculture environment. Competition for resources has been

considered a prevalent force in structuring plant populations under natural selection, often favoring the most competitive individual plants in a particular environment.

The architecture and behavior of successful genotypes as individual plants differ from that of genotypes thriving in a community. Individual plant fitness is increased by 'selfish' traits, which may, like in humans, negatively impact the performance of the community

"Agriculture relies on community performance," emphasizes Prof. Dr. Thorsten Schnurbusch, head of the research group "Plant Architecture" at IPK Leibniz Institute. "But the environment in which crops are grown, i.e. their ecology in the agricultural context, their agroecology, is hardly explored and less understood. It is surprising how less we know about the interactions among plants grown in a dense, real-world community."

Today, crop plants are grown in high-density stands where they experience limited light availability due to mutual shading. "Therefore, by simulating canopy shade, we may get closer to the conditions plants are experiencing in high-density stands in the field, which may be helpful for studying and selecting plants for higher grain yield," says Dr. Guy Golan, first author of the current study. "Cooperative behaviors and highly fertile inflorescences in a light-limited/shaded environment are most important for a thriving grain crop community."

The researchers found behaviors that nourish the fitness of the individual plant as non-beneficial and, in some cases, detrimental to the performance of the whole community. The results have recently been published in the *Plant, Cell & Environment* journal as part of the Special Issue: Tradeoffs in Plant Responses to the Environment.

Moreover, the researchers say that multiple phenotypes attained under simulated shade could better explain community performance of the

wheat crop, advocating the use of simulated shade in breeding high-yielding cultivars.

"Having much deeper insights into these interactions, and specifically understanding their molecular and [genetic components](#) is very important to develop more resilient and resource-efficient [crop plants](#) for the future," says Prof. Dr. Thorsten Schnurbusch. "Embracing an agroecological genetics approach may optimize communal yield by better matching [crops](#) to their environment, as either monoculture or a mixture."

**More information:** Guy Golan et al, Exploring the trade-off between individual fitness and community performance of wheat crops using simulated canopy shade, *Plant, Cell & Environment* (2022). [DOI: 10.1111/pce.14499](#)

Provided by Leibniz Institute of Plant Genetics and Crop Plant Research

Citation: How does the social behavior of wheat plants influence grain production? (2023, September 6) retrieved 12 May 2024 from <https://phys.org/news/2023-09-social-behavior-wheat-grain-production.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------