


# Research team discovers dual effects of chelerythrine in fighting mobile colistin resistance

September 28 2023



(a) Without CHE, the LPS is modified by MCR-1, and colistin is expelled by efflux pumps, where it is difficult for colistin to disrupt the membrane structure. (b) CHE is able to penetrate into the phospholipid bilayers of the plasma membrane and increase the fluidity of the membrane, which inhibits the cellular respiration, disrupts the PMF, and generates ROS, leading to intracellular ATP depletion. (c) Since ATP is critically important, the function of *mcr-1* is limited and the ratio of modified lipid A declines; the function of the efflux pumps is also impaired, which reverses the colistin resistant phenotype. (d) The ATP level is decreased by CHE and the genes involved in the conjugation bioprocess are downregulated, leading to a lower conjugation rate. Credit: Huangwei Song et al.

A research team from China has made an innovative discovery in the fight against mobile colistin resistance. Their study, published in *Engineering*, reveals the dual effects of feed-additive-derived chelerythrine in combating the spread of the *mcr-1* gene, which poses a challenge to the use of colistin, a last-resort antibiotic. This finding opens up new possibilities for enhancing the efficacy of colistin and limiting resistance-gene transmission.

Colistin is a crucial antibiotic used to treat severe infections caused by extensively drug-resistant (XDR) Gram-negative pathogens. However, the emergence and spread of the *mcr-1* gene and its variants have rendered colistin less effective against these pathogens.

In response to this challenge, the research team explored the potential of antibiotic adjuvants to enhance the effectiveness of colistin. Their study focused on the effects of chelerythrine, a compound derived from *Macleaya cordata* extract commonly used as an animal feed additive.

The team discovered that chelerythrine, at a concentration of  $4 \text{ mg}\cdot\text{L}^{-1}$ , significantly reduced the minimal inhibitory concentration (MIC) of colistin against an *mcr-1* positive *E. coli* strain by 16 folds. This reduction is a promising development in combating colistin-resistant pathogens.

In addition, chelerythrine eliminated approximately  $10^4$  colony-forming units (CFUs) of an *mcr-1*-carrying strain in a murine intestinal infection model. Moreover, it inhibited the conjugation of an *mcr-1*-bearing plasmid in vitro by more than 100 folds and in a mouse model by up to 5 folds.

The researchers conducted a detailed analysis to understand the mechanism behind chelerythrine's dual effects. They found that chelerythrine binds to phospholipids on bacterial membranes, increasing

cytoplasmic membrane fluidity.

This alteration impairs respiration, disrupts the proton motive force (PMF), generates [reactive oxygen species](#) (ROS), and decreases intracellular adenosine triphosphate (ATP) levels. As a result, chelerythrine downregulates the *mcr-1* gene and conjugation-associated genes, limiting the spread of mobile colistin resistance.

The implications of this discovery are far-reaching. The dual effects of chelerythrine expand the potential of antibiotic adjuvants, providing a new strategy for combating mobile colistin resistance. This breakthrough may also serve as a reference for future antimicrobial and adjuvant development. By identifying the antimicrobial-resistance-combating effects of known compounds, researchers can explore additional ways to combat the growing threat of antibiotic resistance.

The research team's findings have significant implications for the field of medicine and public health. As antibiotic resistance continues to rise, innovative strategies like chelerythrine's dual effects offer hope in the ongoing battle against drug-resistant pathogens. Further research and development in this area may lead to the discovery of new adjuvants and therapies that can effectively combat mobile [colistin](#) resistance.

The study conducted by the Chinese research team serves as a testament to the importance of interdisciplinary collaboration and the potential of exploring natural compounds for antimicrobial purposes. By harnessing the power of nature and understanding the mechanisms of action, scientists can unlock new strategies to address the global threat of antibiotic [resistance](#).

**More information:** Huangwei Song et al, Dual Effects of Feed-Additive-Derived Chelerythrine in Combating Mobile Colistin Resistance, *Engineering* (2023). [DOI: 10.1016/j.eng.2023.06.012](https://doi.org/10.1016/j.eng.2023.06.012)

## Provided by Engineering

Citation: Research team discovers dual effects of chelerythrine in fighting mobile colistin resistance (2023, September 28) retrieved 2 May 2024 from <https://phys.org/news/2023-09-team-dual-effects-chelerythrine-mobile.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.