

Climate change could jeopardize River Panke restoration successes

October 24 2023



The Panke in Berlin is a typical example of an urban river. Credit: Christian Marx

The Panke River in Berlin is exemplary for many urban watercourses whose water quality has been improved through targeted management.



A team led by the Technical University of Berlin (TU Berlin) and IGB has studied how the <u>water budget</u> and quality have developed over the last 60 years: Restoration measures between 1985 and 1995 have significantly improved the chemical <u>water quality</u>. But <u>climate change</u> could reverse this positive development. Heavy rainfall and dry periods change the chemodynamics of the river.

The Panke is a small river that rises in the Barnim near Bernau and flows into the Spree in Berlin. Of its 29 kilometers of flow, 20 kilometers are on Berlin city territory. "Stinky Panke" it was called in the 19th century, when industry and growing settlements turned the river into a cesspool. But from the 1980s onward, the Panke steadily improved.

"Panke 2015" was the first joint project of the states of Berlin and Brandenburg to implement the EU Water Framework Directive. "Since then, a lot has been invested in renaturation measures, but the Panke is and remains a highly urbanized river that is exposed to diverse human influences and <u>environmental changes</u>," said Dr. Christian Marx, researcher at TU Berlin who conducted the study as part of his doctorate.

The research team evaluated more than 30 years of data on hydroclimate and more than 60 years of data on water quantity and quality of the Panke to find out how the effects of management measures and changing environmental conditions affected the river. The work is <u>published</u> in the journal *Science of the Total Environment*.

"Although the lower reaches of the catchment are still fed by the treated wastewater of about 700,000 people, and about 30% of the catchment is used for agriculture, the water quality in terms of nitrogen nutrients and phosphate is generally good to sufficient for an urban water body and has improved significantly over time at most sites," said Professor Dörthe Tetzlaff, Head of Department at IGB and co-author.



The data show that this is a result of changes in management, restoration and more effective wastewater treatment. The general decrease in fertilizer use has also led to lower nitrate and ammonium pollution. These positive developments were particularly evident after management measures between 1985 and 2000, where investments led to an improvement in water quality from quality Class IV to Class II or better. With Class II status, the German target for meeting the chemical requirement of the European Water Framework Directive has also been achieved in the upper reaches.

Ammonium and phosphate increase in dry periods, nitrate decreases

The long-term data also show that water chemistry has deteriorated again due to hydroclimatic dynamics such as long dry periods, as in 2018, and heavy rainfall events, as in 2017. "In general, the influence of hydroclimate on water quality increases with the degree of urbanization, from upstream to downstream. Hydroclimate also influences the function of water treatment in <u>wastewater treatment plants</u>. So, for a river like the Panke, which is largely fed by treated wastewater in dry periods, this can play a significant role," explained Marx.

When precipitation is lacking in dry periods, pollutants enter wastewater treatment plants less diluted. Wastewater treatment plants with activated sludge treatment may benefit from lower volumes with longer retention times and higher biological activity during drought periods, but at the same time ammonium concentrations increase, suggesting inhibition of nitrification or a concentration effect. However, the increased ammonium concentrations are—at least so far—harmless due to the generally improved water quality and resilience.

Concentrations for phosphate were highest during extremes: drought and



long wet periods. This is due to altered hydrological connectivity and substance-specific flow pathways. Nitrate concentrations were lower during dry periods and higher during wet periods as a result of differences in the mobility and connectivity of agricultural soils to the stream network upstream in the catchment.

However, impacts on water quality classes below Class II were not exceeded. "These findings from the Panke catchment underline the importance of strategic adaptation and improvement of water treatment and water resource management to improve the quality of urban watercourses. They also show how important long-term, integrated data sets are. That is why we are also continuing our detailed process-based research and investigations on the Panke," concluded Tetzlaff.

More information: C. Marx et al, Effects of 66 years of water management and hydroclimatic change on the urban hydrology and water quality of the Panke catchment, Berlin, Germany, *Science of The Total Environment* (2023). DOI: 10.1016/j.scitotenv.2023.165764

Provided by Forschungsverbund Berlin e.V. (FVB)

Citation: Climate change could jeopardize River Panke restoration successes (2023, October 24) retrieved 12 May 2024 from <u>https://phys.org/news/2023-10-climate-jeopardize-river-panke-successes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.