


Decoding the 'montmorency' sour cherry genome: Unraveling prunus evolution and enhancing breeding strategies

October 30 2023



Syntenic relationships between "Montmorency" subgenomes A, A," and B and other *Prunus* species. Credit: *Horticulture Research*

The sour cherry (*Prunus cerasus* L.) is a treasured temperate fruit, known for its distinct sweet-acidic flavor and exceptional processing quality. Belonging to the economically vital Rosaceae family, it shares lineage with various *Prunus* species such as peach, sweet cherry, and almond.

Tracing the evolutionary journey of *Prunus* has historically been challenging, given the widespread hybridization and polyploidy. Studies have identified sour cherry as an allotetraploid ($2n = 4x = 32$), a result of a hybridization between *Prunus fruticosa* Pall. and *Prunus avium* L. Furthermore, climatic challenges like early spring freezes pose significant threats to the temperate fruit industry, causing massive crop loss.

Addressing this, Michigan State University's sour cherry breeding program aims to comprehend the genetic control of bloom time. However, the absence of a public sour cherry genome sequence has limited these studies to traditional linkage maps and marker methods. Therefore, developing a sour cherry genome resource is essential for supporting breeding efforts, enabling [gene discovery](#), and understanding the genetic foundation of agronomic traits in this complex tetraploid.

In May 2023, *Horticulture Research* [published](#) a [research paper](#) entitled "Genome of tetraploid sour cherry (*Prunus cerasus* L.) 'Montmorency' identifies three distinct ancestral *Prunus* genomes."

Researchers first assembled the sour cherry "Montmorency" genome using a combination of PacBio and Illumina sequencing, further scaffolded with chromosome conformation capture (Hi-C). Although expecting two full haplotypes due to the cherry's allotetraploid nature, they discovered 24 linkage groups. Eight of these hinted at origins from the *P. avium*-like ancestor.

Further analyses, including k-mer clustering, segregated these chromosomes into three distinct sets named A, A', and B, representing different subgenomes derived from separate *Prunus* species. An examination of the Illumina reads revealed that subgenome B had twice the genome dosage compared to A or A', confirming the "Montmorency" genome structure as AA'BB.

For structural annotation of gene space, the researchers predicted 92,783 protein-coding genes using RNA sequencing and long-read cDNA datasets from the "Montmorency" tissue. With further analysis using defusion, they found 906 potentially fused genes in the full assembly, of which 707 were present in the scaffolded assembly.

Then, 4,481 [genomic regions](#) on scaffold components were locally relabeled using MAKER within defusion, and 9,777 defused gene models obtained through another Pfam domain search were also re-added to the final annotation. The annotated genes had high BUSCO completion scores and displayed an AED distribution suggesting a high-quality annotation.

For *Prunus fruticosa*, a probable allotetraploid, a draft genome was assembled using PacBio and Illumina reads. Analysis indicated that *P. fruticosa* is also an allotetraploid, with its assembly showcasing common issues linked to polyploid assemblies. Its annotation predicted 102,361 [protein-coding genes](#). Progenitor assignment indicated "Montmorency" subgenomes A and A' as *P. fruticosa*-like, while subgenome B was *P. avium*-like.

In addition, the Dormancy-Associated MADS-box genes (DAMs), essential for *Prunus* dormancy and blooming, were analyzed in "Montmorency" and *P. fruticosa*. Using BLAST+ and manual adjustments via Apollo v 2.6.5, researchers identified three full DAM1–DAM6 haplotypes in "Montmorency."

One, DAM6 on chr1A', had a unique intron-exon structure. *P. fruticosa* exhibited two contigs with full DAM haplotypes, mirroring those in "Montmorency." Additionally, four S-alleles, associated with *Prunus* self-incompatibility, were detected in "Montmorency" with over 99% sequence resemblance to prior studies.

P. fruticosa revealed an S-allele akin to *P. cerasus* S_{36b}, suggesting "Montmorency" sour cherries derived their S₃₆ from a *P. fruticosa* ancestor. The research also indicated that *P. cerasus* "Montmorency" evolved from a hybridization event less than 1.61 million years ago.

In conclusion, this study presents a detailed chromosome-scale genome assembly for the predominant U.S. sour cherry cultivar, "Montmorency," revealing its trigenomic composition of AA'BB. Additionally, the genome of *P. fruticosa*, a key ancestor, was assembled and utilized alongside a *P. avium* sequence.

Key gene sets, including the Dormancy-Associated MADS-box genes (DAMs) and self-incompatibility S-alleles, were annotated. This research enriches our understanding of *Prunus* evolution and offers crucial insights for breeding and comparative genomics, promising advancements in sour cherry cultivation and addressing issues like low fruit set.

More information: Charity Z Goeckeritz et al, Genome of tetraploid sour cherry (*Prunus cerasus* L.) 'Montmorency' identifies three distinct ancestral *Prunus* genomes, *Horticulture Research* (2023). [DOI: 10.1093/hr/uhad097](https://doi.org/10.1093/hr/uhad097)

Provided by NanJing Agricultural University

Citation: Decoding the 'montmorency' sour cherry genome: Unraveling prunus evolution and enhancing breeding strategies (2023, October 30) retrieved 2 May 2024 from <https://phys.org/news/2023-10-decoding-montmorency-sour-cherry-genome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.