

# **Elephant teeth: How they evolved to cope** with climate change-driven dietary shifts

October 16 2023, by Steven Zhang



Credit: Unsplash/CC0 Public Domain

Seeing elephants in the wild is a timelessly awe-inspiring experience.



There are only three living species today: the African savannah elephant, African forest elephant, and Asian elephant.

They are the remnants of a once prosperous lineage of megaherbivores called proboscideans, whose evolutionary epic <u>spanned 60 million years</u> and some 200 species. The African continent was the center stage of this story.

But, until now, it hasn't been clear how <u>natural selection</u> favored modern elephants as the only tiny surviving branch of proboscideans.

A <u>new study</u> by paleontologists Juha Saarinen and Adrian Lister provides the answers. Lister supervised my Ph.D., which focused on evolutionary relationships of fossil elephants. Saarinen hosts my current postdoctoral position at the University of Helsinki, as we're pursuing research stemming from these findings.

They focused on the emergence of elephants' highly specialized, multiplated cheek <u>teeth</u> from the primitive dentition of earlier proboscideans. Their conclusions demonstrate that behavioral adjustments in response to changing environments can produce sustained transformative trends in animals' adaptive structures over tens of millions of years.

# **Teething problems**

Modern elephants' diets consist of copious amounts of tough, fibrous, often low-quality fodder. They're able to eat this way because of their teeth: six massive grinding teeth per jaw half that erupt and expend sequentially in a <u>horizontal "queue"</u> throughout the animal's life, each successively larger than the previous tooth. The adult molars are as big as bricks.

Enamel ridges that align longitudinally along the tooth shear up the



vegetation like a cheese grater as the jaws move. These teeth also have high crowns that continuously erupt into position as the older chewing surface wears away, like lead pushing up in a mechanical pencil. This helps elephants deal with continuous tooth attrition from chewing.

The <u>fossil record</u> shows that, 20 million years ago, proboscidean teeth looked completely different. These teeth had rounded cusps aligned in pairs, versatile for chewing up a wide range of plant food in the primeval forests and thickets they inhabited. But they were ill-suited for tackling large quantities of tough, abrasive vegetation.

Scientists theorized that two major factors drove the evolution of wearresistant dental features seen in elephants today. One was the spread of grass that could tolerate a cooler, drier and more seasonal climate than the cozy forest homes ancestral proboscideans occupied. The second was that <u>large herbivores</u> ingested more adhesive dust in these new arid environments than they had before.

Both challenges arose within the past 20 million years. But it wasn't clear exactly how these two factors drove <u>adaptive changes</u> in herbivore teeth. The theory also presented a chicken-and-egg paradox: which took off first, the evolving teeth or the altered habitats?

## The evidence

Saarinen devised a simple quantitative method to solve this conundrum. It's founded on a basic mechanical principle. The way grazers eat wears the teeth flat, while browsers' teeth have a more "edged" chewing surface. That means grazing elephants should develop angles of wear on their teeth that are wider than those of browsing elephants.

To test this idea, Saarinen first measured recent elephant teeth from Kenya's Tsavo East National Park, which <u>confirmed his initial</u>



postulation. He and Lister then turned to Africa's wealth of geological and fossil records.

Their tooth wear measurements from fossils revealed that, about 21 million years ago, the first increase in dietary uptake of grass was achieved by the primitive "gomphothere"-type proboscideans of Africa. Gomphotheres were the generalized ancestral stock from which modern elephants and their other extinct relatives evolved.

They also discovered that these gomphotheres grazed more as grass cover expanded across their habitats between 21 million and 6 million years ago, as indicated by plant fossil fragments from the sites where the gomphothere teeth were found. Yet this was achieved with little structural evolution in their teeth.

This was a boom period for African proboscidean diversity. Habitats containing three or four proboscidean species were common. Some species adopted grazing to minimize competition with neighboring pachyderms that preferred softer fodder.

### **Ultimate survivors**

To scrutinize previously theorized links between climate, vegetation change and elephant evolution, Saarinen and Lister examined records from <u>sediment cores</u> drilled off Africa's west and east coasts over previous decades. Those cores are essentially archives, capturing tens of millions of years of environmental changes on the continent. Arid periods are marked by dustier segments of the core.

Statistical testing of the data showed that the African environment began to get drastically more arid between 7 million and 5 million years ago, and this drove a vast increase in the rate of dental evolution in the lineage that gave rise to mammoths and modern elephants, as well as



promoting the further spread of grasslands.

Thus came proof that the high-crowned, multi-plated molars of elephants evolved primarily to withstand grit, rather than fiber, in their food. This new dental form provided such an advantage that, <u>around 3.6</u> <u>million years ago</u>, early elephants had outcompeted the last African gomphotheres into extinction.

### New research avenues

The zenith of this evolutionary ratchet was <u>Palaeoloxodon</u>. It was the <u>largest elephant to evolve</u>, standing over 4 meters tall at the shoulder and weighing 12–15 tons. With brick-sized molars packing up to 21 grating plates, Palaeoloxodon was the undisputed top megaherbivore of the African savannah between 1.5 million and around 130,000 years ago, in both its stature and feeding adaptations.

Yet, within the last million years of our story, when African savannahs became <u>increasingly arid and seasonal</u> just as they did for the previous 5 million years, these spectacular grazing specialists vanished. Why?

Upon discussing his findings, Saarinen and I converged on a hypothesis. Modern African elephants are half the size of their prehistoric cousins, with teeth less specialized towards a grass-dominated diet. This may be key for <u>elephants</u> to tolerate the arid shrublands of today's Africa, which have little grass cover to support such enormous grass-dependent vegetarians as Palaeoloxodon.

We're examining this as part of our current research, hoping to decipher whether the encroachment of grasslands by shrubby arid zones drove the extinction of Palaeoloxodon and other spectacular giant herbivores in Africa over the past million years. The story we ultimately tell may serve as a stark warning for how climate change will further decimate African



biodiversity, if it's left unchecked.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Elephant teeth: How they evolved to cope with climate change-driven dietary shifts (2023, October 16) retrieved 9 May 2024 from <u>https://phys.org/news/2023-10-elephant-teeth-evolved-cope-climate.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.