


Study investigates impact of extreme weather events on ocean circulation in tropical Pacific

October 13 2023



ASV change, sea surface temperature, and mixed layer depth. **a** Mixed layer depth (MLD) difference (percent) between GW-ASVCTL and GW (GW-ASVCTL minus GW). The ASV kinetic energy (KE) difference is represented in black contour. The average GW MLD (m) is represented in red contour. **b** Sea surface temperature (SST) difference (°C) between GW-ASVCTL and GW. The ASV kinetic energy difference is represented in black contour. The average SST (°C) is represented in red contour. **c** Mixed layer depth (MLD) difference (percent) between GW-ASVCTL-FLX and GW. The MLD difference (percent)

between GW-ASVCTL and GW is represented in contour. **d** Average SST difference ($^{\circ}\text{C}$) between GW-ASVCTL-FLX and GW. The SST difference ($^{\circ}\text{C}$) between GW-ASVCTL and GW is represented in contour. **e, f** The region 10°N – 10°S (**e**) and 20°S – 40°S (**f**) are considered. The mean MLD (m; abscissa) vs. KE ASV ($\text{m}^2 \text{s}^{-2}$; ordinate) vs. SST anomaly compared to GW-ASVCTL (color) for each experiment are represented. The black circle represents the experiment CTL, the black square GW, the green circle GW-ASVCTL, the green square GW-ASVCTL-FLX (the ASV value corresponding to GW-ASVCTL is plotted here), the blue triangle GW-ASVM10, the blue hexagon GW-ASVM20, the red triangle GW-ASVP10, and the red hexagon GW-ASVP20. Credit: *npj Climate and Atmospheric Science* (2023). DOI: 10.1038/s41612-023-00459-3

The strength of the wind has an important influence on ocean circulation. This is particularly true for extreme events such as storm fronts, tropical storms and cyclones. These weather patterns, which last from a few days to a few weeks, will change in the future due to climate change. In particular, the average energy input into the ocean from mid-latitude storms is expected to decrease, while equatorial regions will become more active. Scientists call these different weather patterns "Atmospheric Synoptic Variability" (ASV).

Two climate researchers—Dr. Olaf Duteil from the GEOMAR Helmholtz Center for Ocean Research Kiel and Professor Dr. Wonsun Park from the IBS Center for Climate Physics and Pusan National University, Korea—have now for the first time investigated the integrated effects of long-term changes in these [weather patterns](#) on the Pacific basin in a modeling study. The results show how important it is to take these changes into account in [climate models](#). They have now [published](#) their findings in the journal *npj Climate and Atmospheric Science*.

From a climate point of view, the weather is usually considered as "noise" and is not systematically analyzed in long-term climate projections, say the two researchers. "However, it is not enough to look at average atmospheric properties, such as mean wind speeds, to understand the influence of [climate change](#) on the ocean," says Duteil, "it is crucial to consider the cumulative effect of short-term changes in weather patterns to get a complete picture."

The researchers expect that future changes in Atmospheric Synoptic Variability will affect the mixing of the ocean's layers, as a smaller or larger input of kinetic energy into the ocean due to weather phenomena will lead to less or more mixing, respectively. The researchers predict that the reduction in ASV in [subtropical regions](#) will lead to a shallowing of the mixing layer in the ocean, while it will become deeper at the equator as ASV increases

They also show that a future reduction in ASV decreases the strength of oceanic circulation systems—the so-called subtropical and tropical cells—and the large-scale [ocean circulation](#). These systems connect mid-latitudes and equatorial latitudes via upper ocean pathways. They are driven by the trade winds north and south of the equator, regulate the upwelling of equatorial waters and play a fundamental role in determining the surface temperature of the oceans and thus primary productivity in the tropics.

This study highlights the need to better quantify ASV and weather patterns in climate models, as changes in ASV have a large impact on future upper ocean circulation and mean properties. Duteil states, "This quantification should be used to improve our confidence in projections of future climate, especially when analyzing large ensembles of climate models."

More information: Olaf Duteil et al, Future changes in atmospheric

synoptic variability slow down ocean circulation and decrease primary productivity in the tropical Pacific Ocean, *npj Climate and Atmospheric Science* (2023). [DOI: 10.1038/s41612-023-00459-3](https://doi.org/10.1038/s41612-023-00459-3)

Provided by Helmholtz Association of German Research Centres

Citation: Study investigates impact of extreme weather events on ocean circulation in tropical Pacific (2023, October 13) retrieved 27 April 2024 from <https://phys.org/news/2023-10-impact-extreme-weather-events-ocean.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.