


Little-known beaked whale found in cold waters off Hokkaido

October 5 2023



Left side full-body photo of the ginkgo-toothed beaked whale described in this study. Credit: Wojtek Bachara, Kuroda Mika, et al. *Aquatic Mammals*, July 9, 2023

Cetaceans are the best-known fully aquatic animals, including whales, dolphins and porpoises. The group includes more than 90 existing

species, divided into baleen whales (Mysticeti) and toothed whales (Odontoceti). Toothed whales are so named because all species in this family possess teeth. Among them, the distribution and ecology of genus *Mesoplodon* are little known, because they typically live offshore in the ocean and rarely surface.

A team of researchers, including Hokkaido University's Assistant Professor Kuroda Mika at the Field Science Center for Northern Biosphere and Professor Matsuishi Takashi Fritz at the Faculty of Fisheries Sciences, recently reported the discovery of a stranded ginkgo-toothed beaked whale from the coast of Yakumo, Southern Hokkaido. [Their findings](#) were published in the journal *Aquatic Mammals*.

"The genus *Mesoplodon* consists of over 15 known species, and is the largest genus of Ziphiidae," Kuroda explains. "The different species in this genus can be identified by the shape of head and by specific teeth in the males. Males of the ginkgo-toothed beaked whale, *Mesoplodon ginkgodens*, possess 10-cm-wide [teeth](#) shaped like the leaves of the ginkgo tree, *Ginkgo biloba*."

Everything we know about ginkgo-toothed beaked whales comes almost exclusively from 95 individuals from 88 separate whale stranding events. Of these, 30 stranding events occurred across Japan.

On February 4, 2022, a whale beaching in Yakumo town, Hokkaido, was reported. The dead whale body was transported to Hakodate Research Center for Fisheries and Oceans for measurement and a necropsy. The whale was a male measuring 477 cm in [body length](#), and was in early stages of decomposition, indicating it had been dead for some time. Its morphology was consistent with those for *M. ginkgodens*; furthermore, genetic analysis of mitochondrial DNA placed this specimen within the clade for *M. ginkgodens*, with one identical sequence.

Portion of the phylogenetic tree showing that the whale in this study (SNH22005) is placed in the *Mesoplodon ginkgodens* clade.

Previous strandings have occurred in a wide range of locations, including Japan, the U.S. West Coast, Australia, the Galapagos Islands, Thailand, New Zealand, Micronesia, the Marshall Islands, China, South Korea and Mexico—all in temperate, subtropical, and tropical waters. This study is the first report of *Mesoplodon ginkgodens* from the colder waters of the North Pacific.

"Another stranding that might have been a ginkgo-toothed beaked whale was reported on November 29, 2021, but the specimen was lost due to bad weather," said Kuroda. "Our findings indicate that these [whales](#) may have migrated near Hokkaido during the winter."

More information: Wojtek Bachara et al, Northernmost Record of the Ginkgo-Toothed Beaked Whale (*Mesoplodon ginkgodens*), *Aquatic Mammals* (2023). [DOI: 10.1578/AM.49.4.2023.356](https://doi.org/10.1578/AM.49.4.2023.356)

Provided by Hokkaido University

Citation: Little-known beaked whale found in cold waters off Hokkaido (2023, October 5) retrieved 21 July 2024 from <https://phys.org/news/2023-10-little-known-beaked-whale-cold-hokkaido.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.