


Examining how Mei-yu precipitation responds to climate change

October 26 2023



Spatial distribution of long-term trends in (a) NDWOR (unit: d/10a), (b) intensity of rainfall event (unit: (mm d⁻¹)/10a), (c) frequency of EPE (unit:

d/10a), (d) intensity of EPE (unit: $(\text{mm d}^{-1})/10\text{a}$), (e) surface air temperature (unit: $^{\circ}\text{C}/10\text{a}$), (f) surface q_s (unit: $(\text{g kg}^{-1})/10\text{a}$), (g) total column water vapor (unit: $\text{kg}/10\text{a}$), and (h) RH (unit: $\%/10\text{a}$) during the Mei-yu period during (a) 1961–2012, (b–e, g) 1961–2022, and (f, h) 1961–2020. The “+” symbols in (a–f) and (h) and the dots in (g) denote where the trend is significant at the 90% confidence level based on the Student’s t-test. The station data in Taiwan Province is not obtained. Credit: Science China Press


Mei-yu (i.e., plume rain) is a distinct weather phenomenon in East Asia during summer, which is generally characterized by persistent rainy and cloudy weather in the middle-lower Yangtze River valley (hereinafter referred to as YRV) region from mid-June to early July. Under global warming, anomalous Mei-yu has caused severe meteorological disasters during recent years.

For instance, in 2020, extreme precipitation events frequently occurred in the YRV region during the Mei-yu period, which caused flood and resulted in over 200 deaths/missing persons and over 170 billion CNY of direct economic losses. In 2022, persistent high temperature and drought events occurred in the YRV region during the Mei-yu period, which greatly affected the agriculture, hydropower, and human health. These [extreme events](#) during the Mei-yu period have brought severe challenges to the government for combating [climate change](#).

[Published](#) in the journal *National Science Review*, the study was led by Prof. Huijun Wang (Nanjing University of Information Science & Technology) and Prof. Bo Sun (Nanjing University of Information Science & Technology), and investigates the changes in the characteristics of Mei-yu under [global warming](#) and the potential reasons based on observation and reanalysis data during 1961–2022.

"The number of days without rainfall (NDWOR), intensity of rainfall

event, and frequency and intensity of extreme precipitation events (EPE) in the YRV region during the Mei-yu period (June 15–July 10) show increasing trends over the past six decades," Prof. Huijun Wang says. "These trends indicate that the weather during the Mei-yu period is becoming more unstable and extreme under global [warming](#)."




(a) Time series of areal mean intensity of regional rainfall events in the YRV region during the Mei-yu period during 1961–2022 (units: mm d⁻¹). Composite differences in transient (b) vertically integrated water vapor flux (unit: 10⁻⁵ kg m⁻¹ s⁻¹; vector) and water vapor divergence (unit: kg m⁻² s⁻¹; color), (c) 850-hPa wind (unit: m s⁻¹; vector) and associated divergence (unit: 10⁻⁶ s⁻¹; color), and (d) 500-hPa vertical velocity (unit: Pa s⁻¹) associated with regional rainfall events in the YRV region during the Mei-yu period between P1 and P2 (P2 minus P1). Composite differences in the (e) intensity of regional rainfall events (unit: mm d⁻¹) in the YRV region during the Mei-yu period between P1 and P2 (P2 minus

P1). The red and blue lines in (a) denote the long-term trend and nine-year sliding average time series, respectively. The yellow dots in (b), (c), (d), the vectors in green in (b), (c), and the “+” symbols in (e) denote where the differences are significant at the 90% confidence level based on the Student’s t-test. The station data in Taiwan Province is not obtained in (e). Credit: Science China Press

The team found that the increasing trend of NDWOR during the Mei-yu period is related to decreased near-surface [relative humidity](#). According to the Clausius-Clapeyron equation, the saturation specific humidity (q_s) would dramatically increase as the global warming continues, at an increasing rate of approximately 7% per °C rise in temperature. As q_s increases more dramatically than q over the YRV region under global warming, the RH is decreased, which may lead to more days without rainfall during the Mei-yu period.

Another possible reason for the increased NDWOR is that the rainfall event may remove $\geq 7\%$ more moisture from the air per °C of local warming. Hence, the air may need a longer time to be replenished with [water vapor](#) before next rainfall event in a warming climate.

Prof. Bo Sun pointed out that the increased intensity of rainfall event and frequency/intensity of EPE during the Mei-yu period may be attributed to combined effects of thermodynamic and dynamic factors. "In a warming climate, the regional rainfall events during the Mei-yu period are associated with stronger water vapor transport, convergence, and convections in the troposphere, which result in increased intensity of rainfall events and hence increased frequency of EPE," Bo Says.



(a) Time series of areal mean surface temperature anomalies (relative to pre-industrial levels during 1861–1900) for individual models and MME under (a) SSP1-2.6 and (e) SSP5-8.5 scenarios. Projected changes (relative to 1985–2005) of (b, f) NDWOR (unit: d), (c, g) intensity of rainfall event (unit: mm d⁻¹) and (d, h) frequency of EPE (unit: d) during the Mei-yu period under 2°C warming scenario based on (b, c, d) SSP1-2.6 and (f, g, h) SSP5-8.5 experiments. The gray dots in (b), (c), (d), (f), (g), (h) denote where the differences are significant at the 90% confidence level based on the Student’s t-test. The red pentacle symbols and horizontal gray dashed lines in (a) and (e) denote where the temperature rising reaches 2°C of global warming relative to pre-industrial climate. Credit: Science China Press

Furthermore, the researchers analyzed the response of Mei-yu to 2°C of global warming with respect to pre-industrial climate using CMIP6 models. The results suggest that the NDWOR, intensity of [rainfall](#)

events, and frequency of EPE will be increased in the YRV region during the Mei-yu [period](#) under the 2°C warming scenario, which imply a more challenging climate risk management in the future.

More information: Bo Sun et al, How does Mei-yu precipitation respond to climate change?, *National Science Review* (2023). [DOI: 10.1093/nsr/nwad246](#)

Provided by Science China Press

Citation: Examining how Mei-yu precipitation responds to climate change (2023, October 26) retrieved 29 April 2024 from <https://phys.org/news/2023-10-mei-yu-precipitation-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.