


Scientists develop acoustic 3D trapping of microparticles in flowing liquid using circular cavity

October 11 2023, by Zhang Nannan



Schematic diagram of the chip design and its trapping performance on red blood cells and white blood cells. The cells aggregate to the center of the circular cavity within 60 ms. Credit: SIBET

A research team from the Suzhou Institute of Biomedical Engineering and Technology (SIBET) of the Chinese Academy of Sciences has developed an acoustic trapping chip that can provide three-dimensional (3D) trapping of cells in a continuously flowing medium with a circular resonance structure.




The study, titled "Acoustic 3D trapping of microparticles in flowing liquid using circular cavity," has been published in *Sensors and Actuators A: Physical*.

Cell trapping is of great importance in <u>biomedical engineering</u> because it enables the clamping, separation, filtration and agglomeration of cells. Among different trapping approaches, acoustic trapping has been widely used in <u>biological research</u> because it can provide contactless and biosafe cell manipulation.

Ultrasonic standing waves can be further categorized into standing bulk acoustic waves (BAW), generated by a bulk <u>piezoelectric transducer</u>, or standing surface <u>acoustic waves</u> (SAW), generated by single crystal lithium niobate (LiNbO₃) etched with interdigitated electrodes. SAW can manipulate particles with very low energy consumption, but it is generally used for sorting in flowing liquid and particle arrangement in stationary liquid due to its overall smaller clamping <u>force</u> compared with BAW.

On the other hand, the acoustic microstreaming vortex can also be applied to trap cells near the obstacle or microbubbles. The design of micropillars or obstacles plays an important role in improving the trapping efficiency. However, some of the traps cannot release particles easily, and some of them cannot provide a fixed trapping position.





Single particle with diameter of $10 \ \mu m$ queuing up on the cavity array was delivered step by step to the detection spot, which can be applied to high throughput Reman spectra acquisition. Credit: SIBET

The trapping efficiency is basically determined by the trapping force. In most previous studies, particles are usually trapped in a static fluid or a fluid flowing at extremely low speed, or the trapping process takes several seconds, which is mainly due to insufficient trapping force. This reduces both trapping efficiency and throughput, while high-throughput cell manipulation is important in many biological applications, such as Raman identification and nanoparticle capture.

The researchers established a standing acoustic wave in the circular microstructure, providing sufficient force to clamp cells in the center of the chamber. Meanwhile, <u>cells</u> near the bottom of the micro channel are clamped under the radiation force generated in the depth direction.



Thus, a 3D cell confinement is formed with a special design of microchannels actuated by only one piezoelectric plate transducer. Experimental results show that the chip can provide nanonewton (nN) level trapping force and millisecond (ms) level trapping time for micron-sized particles moving at the speed of mm/s level.

With this non-contact and biocompatible trapping method, the chip can be applied to a variety of biomedical engineering scenarios such as organ chips, cell culture, Raman analysis and nanoparticle capture.

More information: Ce Wang et al, Acoustic 3D trapping of microparticles in flowing liquid using circular cavity, *Sensors and Actuators A: Physical* (2023). DOI: 10.1016/j.sna.2023.114698

Provided by Chinese Academy of Sciences

Citation: Scientists develop acoustic 3D trapping of microparticles in flowing liquid using circular cavity (2023, October 11) retrieved 9 May 2024 from <u>https://phys.org/news/2023-10-scientists-acoustic-3d-microparticles-liquid.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.