


Scientists present the first set of global maps showing geographic patterns of beta-diversity in flowering plants

October 16 2023



Geographic patterns of taxonomic (a) and phylogenetic (b) beta-diversity in flowering plant genera in regional floras in the world. Credit: Shenhua Qian

Beta-diversity serves as a crucial metric for gauging shifts in species composition over spatial or temporal scales, bridging the spectrum between localized (alpha) and broader regional (gamma) diversity. In the fields of ecology, biogeography and conservation biology, to elucidate the origins and sustenance of geographic beta-diversity patterns, we need to explore both the taxonomic and phylogenetic beta-diversity at different evolutionary depths.

In an [article](#) published in *Plant Diversity*, using a comprehensive database of plant distributions worldwide, Dr. Hong Qian of Illinois State Museum in the US and Dr. Shenhua Qian of Chongqing University in China explored and mapped geographic patterns of beta-diversity for flowering plant (angiosperm) genera in regional floras across the world.

They analyzed both taxonomic and phylogenetic beta-diversity, including their constituent elements. Moreover, they considered both tip-weighted and basal-weighted phylogenetic beta-diversity in relation to latitude.

Consequently, they showed that the global distribution of beta-diversity is highly heterogeneous. This is the case for taxonomic and phylogenetic beta-diversity, as well as for both tip-weighted and basal-weighted phylogenetic beta-diversity. They also showed that there are highly consistent geographic patterns among different metrics of beta-diversity. In most cases, metrics of beta-diversity are negatively associated with latitude, particularly in the Northern Hemisphere.

Notably, different metrics of taxonomic beta-diversity are strongly and positively correlated with their counterparts of phylogenetic beta-diversity.

"It is well known that regions characterized by wet tropical climates tend

to exhibit elevated levels of alpha- and gamma-diversity in both plant and [animal species](#)," says Qian. "However, one of the interesting patterns that we found in this study is that regions of high beta-diversity are located in dry rather than wet tropical climates. For example, the Sahara Desert has the highest beta-[diversity](#) of flowering plants."

More information: Hong Qian et al, Geographic patterns of taxonomic and phylogenetic β -diversity of angiosperm genera in regional floras across the world, *Plant Diversity* (2023). [DOI: 10.1016/j.pld.2023.07.008](#)

Provided by KeAi Communications Co.

Citation: Scientists present the first set of global maps showing geographic patterns of beta-diversity in flowering plants (2023, October 16) retrieved 21 May 2024 from <https://phys.org/news/2023-10-scientists-global-geographic-patterns-beta-diversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
