


# Superatomic semiconductor sets a speed record for ballistic flow

October 26 2023



What makes silicon a desirable semiconductor is that electrons *can* move through it very quickly, but like the proverbial hare, they bounce around too much and don't actually make it very far, very fast in the end. Excitons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  are, comparatively, very slow, but it's precisely because they are so slow that they are able to meet and pair up with equally slow-moving acoustic phonons. The resulting quasiparticles are "heavy" and, like the tortoise, advance slowly but steadily along. Unimpeded by other phonons along the way, acoustic exciton-polarons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  ultimately move faster than electrons in silicon. Credit: Jack Tulyag, Columbia University

Semiconductors—most notably, silicon—underpin the computers,

cellphones, and other electronic devices that power our daily lives, including the device on which you are reading this article.

As ubiquitous as semiconductors have become, they come with limitations. The atomic structure of any material vibrates, which creates [quantum particles](#) called phonons.

Phonons in turn cause the particles—either electrons or [electron-hole pairs](#) called excitons—that carry energy and information around [electronic devices](#) to scatter in a matter of nanometers and femtoseconds. This means that energy is lost in the form of heat, and that information transfer has a speed limit.

The search is on for better options. Writing in *Science*, a team of chemists at Columbia University led by Jack Tulyag, a Ph.D. student working with chemistry professor Milan Delor, describes the fastest and most efficient [semiconductor](#) yet: a superatomic material called  $\text{Re}_6\text{Se}_8\text{Cl}_2$ .

Rather than scattering when they come into contact with phonons, excitons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  actually bind with phonons to create new quasiparticles called acoustic exciton-polarons. Although polarons are found in many materials, those in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  have a special property: they are capable of ballistic, or scatter-free, flow. This ballistic behavior could mean faster and more efficient devices one day.

In experiments run by the team, acoustic exciton-polarons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  moved fast—twice as fast as electrons in silicon—and crossed several microns of the sample in less than a nanosecond.

Given that polarons can last for about 11 nanoseconds, the team thinks the exciton-polarons could cover more than 25 micrometers at a time. And because these quasiparticles are controlled by light rather than an

electrical current and gating, processing speeds in theoretical devices have the potential to reach femtoseconds—six orders of magnitude faster than the nanoseconds achievable in current Gigahertz electronics—all at room temperature.

"In terms of energy transport,  $\text{Re}_6\text{Se}_8\text{Cl}_2$  is the best semiconductor that we know of, at least so far," Delor said.

## **A quantum version of the tortoise and the hare**

$\text{Re}_6\text{Se}_8\text{Cl}_2$  is a superatomic semiconductor created in the lab of collaborator Xavier Roy. Superatoms are clusters of atoms bound together that behave like one big atom, but with different properties than the elements used to build them. Synthesizing superatoms is a specialty of the Roy lab, and they are a main focus of Columbia's Material Research Science and Engineering Center on Precision Assembled Quantum Materials.

Delor is interested in controlling and manipulating the transport of energy through superatoms and other unique materials developed at Columbia. To do this, the team builds super-resolution imaging tools that can capture particles moving at ultrasmall, ultrafast scales.

When Tulyag first brought  $\text{Re}_6\text{Se}_8\text{Cl}_2$  into the lab, it wasn't to search for a new and improved semiconductor—it was to test the resolution of the lab's microscopes with a material that, in principle, shouldn't have conducted much of anything. "It was the opposite of what we expected," said Delor. "Instead of the slow movement we expected, we saw the fastest thing we've ever seen."

Tulyag and his peers in the Delor group spent the next two years working to pinpoint why  $\text{Re}_6\text{Se}_8\text{Cl}_2$  showed such remarkable behavior, including developing an advanced microscope with extreme spatial and temporal

resolution that can directly image polarons as they form and move through the material. Theoretical chemist Petra Shih, a Ph.D. student working in Timothy Berkelbach's group, also developed a quantum mechanical model that provides an explanation for the observations.

The new quasiparticles are fast, but, counterintuitively, they accomplish that speed by pacing themselves—a bit like the story of the tortoise and the hare, Delor explained. What makes silicon a desirable semiconductor is that electrons can move through it very quickly, but like the proverbial hare, they bounce around too much and don't actually make it very far, very fast in the end.

Excitons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  are, comparatively, very slow, but it's precisely because they are so slow that they are able to meet and pair up with equally slow-moving acoustic phonons. The resulting quasiparticles are "heavy" and, like the tortoise, advance slowly but steadily along. Unimpeded by other phonons along the way, acoustic exciton-polarons in  $\text{Re}_6\text{Se}_8\text{Cl}_2$  ultimately move faster than electrons in silicon.

## **The semiconductor search continues**

Like many of the emerging quantum materials being explored at Columbia,  $\text{Re}_6\text{Se}_8\text{Cl}_2$  can be peeled into atom-thin sheets, a feature that means they can potentially be combined with other similar materials in the search for additional unique properties.  $\text{Re}_6\text{Se}_8\text{Cl}_2$  however, is unlikely to ever make its way into a commercial product—the first element in the molecule, Rhenium, is one of the rarest on Earth and extremely expensive as a result.

But with the new theory from the Berkelbach group in hand along with the advanced imaging technique that Tulyag and the Delor group developed to directly track the formation and movement of polarons in the first place, the team is ready to see if there are other superatomic

contenders capable of beating  $\text{Re}_6\text{Se}_8\text{Cl}_2$ 's speed record.

"This is the only material that anyone has seen sustained room-temperature ballistic exciton transport in. But we can now start to predict what other materials might be capable of this behavior that we just haven't considered before," said Delor. "There is a whole family of superatomic and other 2-D semiconductor materials out there with properties favorable for acoustic polaron formation."

**More information:** Jakhangirkhodja A. Tulyagankhodjaev et al, Room-temperature wavelike exciton transport in a van der Waals superatomic semiconductor, *Science* (2023). [DOI: 10.1126/science.adf2698](https://doi.org/10.1126/science.adf2698).  
[www.science.org/doi/10.1126/science.adf2698](https://www.science.org/doi/10.1126/science.adf2698)

Provided by Columbia University

Citation: Superatomic semiconductor sets a speed record for ballistic flow (2023, October 26) retrieved 29 April 2024 from <https://phys.org/news/2023-10-superatomic-semiconductor-ballistic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.