

New flood modeling research aims to produce more accurate predictions

November 1 2023, by Betsy Sonewald





Study area map showing the computational model domain. A solid dark rectangle in the middle depicts the region affected the most by the August 2021 extreme flooding event including Waverly, TN. Credit: *Journal of Hydrology* (2023). DOI: 10.1016/j.jhydrol.2023.130157

On Aug. 21, 2021, Waverly, Tennessee, received nearly one-third of its average annual rainfall in just a few hours—the resulting 17 inches of rain led to one of the deadliest flooding events on record.

In fact, Waverly faced some of the worst consequences of the severe rain and subsequent flooding that hit Central Tennessee in the summer of 2021. Unfortunately, the area was unprepared for the torrent that killed 21 people and caused over \$100 million in property damage.

A combination of factors causes flooding, including rainfall, <u>geographic</u> <u>location</u>, the amount of moisture already in the environment, the flow of rivers and streams and surface runoff. And each factor has a direct impact on the severity of the flood. A variety of models can simulate these individually, but few studies have assessed the accuracy of a framework that captures the entire flood event.

To better understand the factors that contributed to the August 2021 floods, a multi-institutional team tested a new hierarchical flood modeling framework that can simulate the entire flood event and translate precipitation to inundation. One of the main questions with a multimodel approach is how errors propagate through coupled, or dependent, simulations.

In a study <u>published in the *Journal of Hydrology*</u>, researchers identified one of the most important sources of uncertainty, or error, in the flood models and demonstrated the accuracy and utility of the framework for



predicting future floods.

The recent study explored a new approach for flood modeling that could improve predictions for future weather events.

The study was led by Sudershan Gangrade, Ganesh Ghimire and Shih-Chieh Kao from Oak Ridge National Laboratory's Water Resource Science and Engineering group and Mario Morales Hernandez from the University of Zaragoza in Spain.

"We developed a modeling framework to unravel the August 2021 extreme flooding event at Waverly," said Gangrade. "Developing and testing the framework for Waverly is not only useful for retrospective analysis but can also be utilized to simulate future flood scenarios."

In this study, the research team used <u>precipitation data</u> to run two hydrologic models—the Variable Infiltration Capacity model and the Routing Application for Parallel Computation of Discharge, or RAPID—to simulate runoff and streamflow in the Waverly area.

The outputs from those models were then fed into two inundation models—AutoRoute and the Two-dimensional Runoff Inundation Toolkit for Operational Needs, or TRITON, which is a GPU-optimized model designed to run on modern supercomputers.

The team wanted to know how inaccuracies in each model affected the final inundation models, which could be used to inform evacuation orders and emergency response.

To test the impact of different precipitation inputs on the final flood prediction, the team used both forecast data and observed data from the Waverly area. Running a model with data from a past weather event is a common practice known as hindcasting, which tests a model's accuracy



compared to observations.

"We wanted to perform a diagnostic evaluation of the Waverly flood," said Ghimire. "We saw significant differences between the inundation <u>model</u> outputs from the forecast and observed precipitation data. The error in the precipitation forecasts propagated through the framework and was the main reason we underestimated the central Tennessee flood."

Tests using the observed precipitation data showed that the <u>framework</u> accurately simulated the flooding, which led the team to conclude that errors in the inundation forecast primarily originated from the <u>precipitation</u> forecasts.

As forecasts improve and the team works to scale the models to cover larger domains within the U.S. and global watersheds, its usefulness in future flood preparedness and management holds real promise.

The team is excited to explore opportunities to produce more accurate and timely predictions that can help communities better prepare for and respond to <u>flood</u> events.

More information: Sudershan Gangrade et al, Unraveling the 2021 Central Tennessee flood event using a hierarchical multi-model inundation modeling framework, *Journal of Hydrology* (2023). DOI: <u>10.1016/j.jhydrol.2023.130157</u>

Provided by Oak Ridge National Laboratory

Citation: New flood modeling research aims to produce more accurate predictions (2023, November 1) retrieved 12 May 2024 from <u>https://phys.org/news/2023-11-aims-accurate.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.