


# Astronomers discover galaxies have bipolar gas outflows reaching far into intergalactic space

December 6 2023

---




Gas outflows (red) near spiral galaxies (white). The left image shows the outflow of gas for the group of galaxies we see edge on. It is clear that most of the gas escapes upward and downward into intergalactic space. The right image shows the galaxies we see face on. A ring of gas is visible here because the gas in front of the galaxy absorbs more light from the galaxy than it emits itself (net absorption is blue). Credit: HST/ESO/VLT/MUSE/Yucheng Guo et al.

Astronomers have observed, for the first time in three dimensions, that gas from spiral galaxies is blown upward and downward at high velocity, far out of the galaxy. The observations confirm the prevailing theory of galaxy evolution that says that star-forming galaxies drive intergalactic winds by venting their gas along the poles. The astronomers have published their findings in *Nature*.

Gas flows are important in models that describe the formation of galaxies. Galaxies grow by the inflow of gas from their surroundings. Such growth, it is thought, is inhibited by [young stars](#) and [supermassive black holes](#) that blow gas far out into space through [shock waves](#). Exactly what happens is unknown, but without strong gas flows, galaxies would become much too massive.

Researchers have now made the first convincing case that galactic winds reach far beyond galaxies. They used the MUSE instrument on the Very Large Telescope in Chile. They studied the signal from magnesium gas around nearly two hundred, distant, spiral galaxies. In half of these galaxies, they were observing the disk edge-on. In the other half, they saw the disk face-on, as a circle.

In the group of edge-on galaxies, the gas outflow was perpendicularly upward and downward. "We detect the gas up to tens of thousands of [light years](#) from the galaxy, moving through intergalactic space at hundreds of kilometers per second," says first author Yucheng Guo (Université de Lyon, France).



The scientists were able to identify the morphology of galactic winds. In this figure, magnesium atom emissions have been used to trace galactic winds. The flow of matter occurs along the central axis perpendicular to the galactic disc.

Credit: © Yucheng Guo

"To me, it is a real milestone that we are finally witnessing intergalactic gas outflows from ordinary galaxies," says Joop Schaye (Leiden University, the Netherlands). He co-authored the study and has been researching galactic winds and gas flows in intergalactic space for years. "Until now, observations have been difficult to interpret, but thanks to this study we can no longer ignore bipolar winds."

Now that [astronomers](#) have mapped average gas flows and velocities, they can test and adjust their computer simulations of galaxy evolution. This is important because it will clarify how galaxies grow.

**More information:** Yucheng Guo et al, Bipolar outflows out to 10 kpc for massive galaxies at redshift  $z \approx 1$ , *Nature* (2023). [DOI: 10.1038/s41586-023-06718-w](https://doi.org/10.1038/s41586-023-06718-w).  
[www.nature.com/articles/s41586-023-06718-w](https://www.nature.com/articles/s41586-023-06718-w)

Provided by Netherlands Research School for Astronomy

Citation: Astronomers discover galaxies have bipolar gas outflows reaching far into intergalactic space (2023, December 6) retrieved 18 June 2024 from <https://phys.org/news/2023-11-astronomers-galaxies-bipolar-gas-outflows.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.