


# How looking beyond our borders would improve preparedness for extreme floods

November 7 2023



Mega-floods in Europe. **a**, Five hydroclimatic regions: Boreal (purple), Continental (blue), Mediterranean (orange), Alpine (green) and Atlantic (red). **b–f**, Maximum observed specific flood discharges (points) and mean of annual specific flood discharges (squares) over the entire observation period at each stream gauge as a function of catchment area. Regional envelope curves (thick lines) and median regional annual specific flood discharges (thin lines) for the full record period are shown for each hydroclimatic region: Boreal (**b**), Continental (**c**), Mediterranean (**d**), Alpine (**e**) and Atlantic (**f**). Envelope curves for two 30-year sub-periods are also shown (dashed lines for 1961–1990, dotted lines for 1991–2020). Parameters of the envelope curves are listed in Extended Data Table 2. Colored symbols indicate the mean and maximum flood discharges in the 498 catchments with recent mega-floods, gray points those of

the remaining catchments. **g–j**, Examples of series of annual flood discharges with (**g,i**) and without (**h,j**) megafloods; their corresponding mean (squares) and maximum values (points) are highlighted in black in **c** and **f**. The locations of corresponding stream gauges are indicated in **a** by circles. PL, Poland; AT, Austria; ES, Spain; GB, United Kingdom. Credit: *Nature Geoscience* (2023). DOI: 10.1038/s41561-023-01300-5

Almost all extreme flooding across Europe could be anticipated by looking at previous major events in other similar parts of the continent, according to a new study.

"Megafloods," such those on the Rhine tributaries in Germany in 2021, are [extreme events](#) that vastly exceed those previously experienced in each location, causing significant damage and deaths. But they can be hard to anticipate due to their rarity and the lack of data at many sites, which means flood defenses and emergency response plans are often insufficient, exacerbating the impacts on communities.

A team of European scientists, including experts at the UK Centre for Ecology & Hydrology (UKCEH), analyzed river discharges—the volume of water flowing through a channel per second—from 8,000 gauging stations across Europe from 1810–2021 to identify historic megafloods.

They found 95.5% of megafloods could have been anticipated based on previous events at locations elsewhere in the continent with similar climate and variability in terms of how much water that rivers discharge in response to factors such as rainfall and temperature.

UKCEH Principal Hydrologist Jamie Hannaford was a co-author of the study, [published](#) in *Nature Geoscience*. He explains, "Our findings demonstrate that while the most extreme floods shock [local communities](#)

, they are not usually surprising if we take a continental-scale viewpoint."

"In the UK we already look beyond local catchments and consider events from other locations when assessing flood risk, but this is still limited to within our borders. A continental scale approach could give us additional information on our susceptibility to extreme floods. This would provide 'worst case scenarios' to help ensure appropriate flood defense measures and preparations can be implemented, thereby limiting damage when extreme floods do happen."

The [catastrophic flooding](#) in July 2021 at the Rhine tributaries in Germany, and rivers in the Netherlands, Belgium and Luxembourg, was up to four times larger than any event on record in the region. It took residents and authorities by surprise, and caused more than 200 deaths and damage worth US \$40 billion. However, the new study showed that the discharge rate was similar to floods in northern Austria in 2002.

Examples in the UK include the 2009 floods in the Derwent catchment in Cumbria, where the discharge was 58% greater than the second largest event on record there, but not as extreme as flooding in similar catchments in Norway.

Mr. Hannaford and fellow UKCEH hydrologist Steve Turner were among the team of 56 researchers representing institutes in 29 countries, led by the Vienna University of Technology.

A [study in 2019](#) by the same research team found that [flood](#) events are becoming increasingly severe in north-western Europe, with northern England and southern Scotland among the areas most affected, but decreasing in severity in southern and Eastern Europe.

**More information:** Miriam Bertola et al, Megafloods in Europe can be anticipated from observations in hydrologically similar catchments,

*Nature Geoscience* (2023). [DOI: 10.1038/s41561-023-01300-5](https://doi.org/10.1038/s41561-023-01300-5)

Provided by UK Centre for Ecology & Hydrology

Citation: How looking beyond our borders would improve preparedness for extreme floods (2023, November 7) retrieved 11 September 2024 from <https://phys.org/news/2023-11-borders-preparedness-extreme.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.