


# Researchers achieve chemically controlled, reversible magnetic phase transition

November 10 2023, by Li Rui



Schematic illustration of the magnetic phase change in a 2D metal-organic lattice induced by the lactic-lactam tautomerization reaction. Credit: USTC

A research team led by Associate Prof. Li Xingxing and Prof. Yang Jinlong from the University of Science and Technology of China (USTC) of the Chinese Academy of Sciences (CAS) has developed a groundbreaking chemical method for two-dimensional metal-organic lattices.


Their results were [published](#) in *Nano Letters* on Oct. 2nd.

In spintronics, it is paramount to develop an [efficient way](#) to reversibly control the spin order of materials. Though various physical methods have been proposed, chemically achieving this has posed significant challenges.

Researchers proposed the utilization of the well-recognized lactim–lactam tautomerization process to reversibly modulate the magnetic phase transition in two-dimensional (2D) organometallic lattices. This revelation offers novel pathways for controlling the electrical and magnetic characteristics of materials.

The spin state of an organic linkers undergoes a transformation from a singlet state to a doublet state due to the lactim–lactam tautomerization.

Using chemical means to control the spin state of materials has several potential advantages over physical methods. It can be done at [room temperature](#), which makes it more practical for real-world applications. Additionally, chemical reactions can be precisely controlled, enabling more [precise control](#) over the spin state of materials.



a-c) Geometric structure, ground state spin density distribution, and band structure of the lactam-type Cr-pyrazine metal-organic lattice. (d-f) Geometric structure, ground state spin density distribution, and band structure of the lactim-type Cr-pyrazine metal-organic lattice. Credit: *Nano Letters* (2023). DOI: 10.1021/acs.nanolett.3c03060

In their study, the team used the compound called 2D organometallic lattices, which has a unique structure that allows its magnetic phase to be changed using the lactim-lactam tautomerization. Researchers demonstrated that this reaction could be used to reversibly switch the

magnetic state of the material from antiferromagnetic to ferrimagnetic.

The team's findings pave the way for further research in this area. By exploring other [chemical reactions](#) that can influence the spin state of materials, it may be possible to develop even more advanced spintronic devices in the future.

**More information:** Junyao Li et al, Chemically Controlled Reversible Magnetic Phase Transition in Two-Dimensional Organometallic Lattices, *Nano Letters* (2023). [DOI: 10.1021/acs.nanolett.3c03060](https://doi.org/10.1021/acs.nanolett.3c03060)

Provided by University of Science and Technology of China

Citation: Researchers achieve chemically controlled, reversible magnetic phase transition (2023, November 10) retrieved 2 May 2024 from <https://phys.org/news/2023-11-chemically-reversible-magnetic-phase-transition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.