

Climate change is hastening the demise of Pacific Northwest forests

November 16 2023, by NATHAN GILLES, Columbia Insight



Credit: Vlada Karpovich from Pexels

Deep inside a forest in Oregon's Willamette Valley stands a dead "Tree of Life."



Its foliage, normally soft and green, is tough and brown or missing altogether. Nonetheless, the tree's reddish bark, swooping branches and thick, conical base identify it as the Pacific Northwest's iconic western red cedar.

Christine Buhl, a forest health specialist for the Oregon Department of Forestry, plunges a tool called an increment borer into the dead tree's trunk. Twisting the handle of the corkscrew-like borer, Buhl extracts a long, thin sample of the tree's inner growth rings.

The rings become thinner over time, indicating the tree's growth slowed before the tree finally died, a sign that this red cedar, like thousands of others in Oregon and Washington, died from drought.

"That's why it's the canary," says Buhl. "Any tree that's less drought tolerant is going to be the canary in the coal mine. They're going to start bailing (out)."

For thousands of years, people have used red cedar to make everything from canoes to clothing.

Red cedar's many uses have earned the species endearing names, including the "Tree of Life." More recently, scientists have started calling this water-loving relative of redwoods by a less flattering name: "the climate canary."

Last year, Buhl and colleagues <u>reported that red cedars were dying</u> throughout the tree's growing range not because of a fungus or insect attack, but due to the region's "climate change-induced drought."

Red cedars aren't alone.

In recent years, at least 15 native Pacific Northwest tree species have



experienced growth declines and die-offs, 10 of which have been linked to drought and warming temperatures, according to <u>recent studies and</u> <u>reports</u>.

Many researchers, Buhl included, are now arguing that these droughtdriven die-offs are the beginning of a much larger and long-predicted shift in tree growing ranges due to climate change.

Trees, and plants generally, have growing ranges that are largely determined by climate factors, namely moisture and temperature.

For decades, scientists have argued that as atmospheric warming continues, growing ranges in the Northern Hemisphere will shift upslope in elevation and farther north, leaving many trees stranded in a warmer, drier world.

As climate mismatch sets in, trees are expected to die-off and not grow back, according to predictions.

Daniel DePinte, Forest Service aerial survey program manager, suspects range changes are driving "Firmageddon." A term coined by researchers, including DePinte, "Firmageddon" refers to the more than 1,875 square-mile (4,856-square-kilometer) die-off of five fir species in Oregon, Washington and northern California.

"The forests are moving uphill," said DePinte.

DePinte and colleagues first identified and named the massive, droughtdriven fir "mortality event" last year while surveying area forests via airplane.

According to tree-range predictions, climate-induced die-offs are expected to start at the edges of growing ranges, including at lower-



elevation locations that are predicted to become too warm and dry for many species.

DePinte's survey revealed that the largest die-offs associated with Firmageddon are occurring at lower-elevation sites.

Buhl and colleagues found a similar pattern with western red cedar. Mortality was greatest at sites less than about 650 feet (200 meters) in elevation west of the Cascade Range, according to their analysis.

Scientists have also observed a similar pattern for Douglas fir, the region's leading commercial timber species. Douglas fir is currently experiencing a 720-square-mile (1,865-square-kilometer) die-off, the majority in the Klamath Mountains near the southern Oregon cities of Ashland and Medford.

The die-off is limited to the lower elevations but is likely to move uphill as temperatures warm in the coming decades, according to a study in the <u>Journal of Forestry</u>.

"Our analysis concluded that if climate change continues as predicted, we could see increased Douglas fir mortality at higher elevations," said study co-author, David Shaw, a professor and forest health specialist at Oregon State University.

Shaw called the die-off "consistent with predictions for climate change."

But whereas red cedar is believed to be dying from drought alone, the Firmageddon and Douglas fir die-offs have been linked to a combination of drought weakening trees and insect pests moving in for the kill.

"These insects are not normally tree killers," said DePinte. "This is evidence that the forests are reacting to climate change and droughts."



Douglas fir is not considered a true fir and is not officially part of Firmageddon, according to DePinte.

The combination of drought-induced stress and pests, said Patrick Tobin, associate professor of disturbance ecology at the University of Washington, is analogous to a person with a weakened immune system dying from the flu.

"Drought-stress opens a window for biotic agents that might otherwise not be able to overcome a healthy, well-defended tree," said Tobin.

Tobin is a co-author of a 2021 study in the journal Forest Ecology and Management on the widespread decline of big leaf maples in western Washington. Tobin's study was unable to determine whether drought alone or drought in combination with disease-causing fungi was killing the native maples.

As for the climate canary, Buhl believes red cedar is unlikely to disappear from the landscape entirely but probably won't grow back in areas where it's dying off.

"Unless we turn back climate change, there is no reason to hope western red cedar is going to make a comeback," said Buhl.

© 2023 The Associated Press. All rights reserved. This material may not be published, broadcast, rewritten or redistributed without permission.

Citation: Climate change is hastening the demise of Pacific Northwest forests (2023, November 16) retrieved 20 May 2024 from https://phys.org/news/2023-11-climate-hastening-demise-pacific-northwest.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.