


Plants found to recruit distinct chemical activities of coumarins under different soil pHs

November 29 2023



Coumarin biosynthesis in roots responds to environmental pH conditions. Under slightly acidic conditions (low pH), plants produce the superior ferric iron (Fe^{3+}) reductant sideretin, which complements Fe^{3+} reduction by the membrane-bound reductase FRO2. At neutral to alkaline conditions, fraxetin is synthesized to maintain the mobilization of poorly soluble iron sources for further reduction by FRO2. Credit: IPK Leibniz Institute

Plants have two main uptake mechanisms to obtain iron (Fe) from the soils. The type of strategy employed depends on the botanical classification of the plant. In the so-called strategy-I mechanism, plants must first reduce the trivalent iron (Fe^{3+}) into bivalent iron (Fe^{2+}). Only

then can Fe^{2+} be taken up by roots as a free ion.

This strategy is used by non-Poaceae plants such as [oilseed rape](#) and the model plant *Arabidopsis thaliana*. Grasses, which belong to the Poaceae family, employ a so-called strategy II. These plants secrete chelating compounds, which can be reuptaken once chelated with Fe^{3+} . Thus, no reduction step is required for Fe import into root cells.

"Interestingly, some Strategy-I plants also release metabolites into the soil through their roots when they suffer from Fe deficiency. Some of these are coumarins," explains IPK scientist Dr. Ricardo Giehl, co-head of the "Molecular Plant Nutrition" working group. However, the physiological role of these coumarins has not yet been sufficiently clarified.

In their work, the researchers were able to show that the main function of two coumarins released in response to Fe deficiency is largely dependent on the external pH. Under slightly acidic conditions, coumarins, especially sideretin, help to sustain Fe^{3+} reduction.

Here, sideretin works together with the membrane-bound enzyme FERRIC REDUCTION OXIDASE 2 (FRO2) to efficiently Fe^{2+} uptake into the roots.

If the medium is alkaline, the biosynthesis of coumarins is shifted from sideretin to fraxetin, a response that the research team found to depend on the transcription factor MYB72. At alkaline pH, sideretin loses both its ability to reduce and even to solubilize Fe^{3+} from precipitated sources, while fraxetin retains a high Fe^{3+} mobilization capacity under such pH conditions. Therefore, rather than reducing Fe^{3+} directly, the main function of fraxetin is to provide soluble Fe(III)-chelates for FRO2-mediated reduction.

"Our study shows that by adjusting [coumarin](#) biosynthesis, plants recruit specific functions depending on the prevailing pH of the soil," says Dr. Ricardo Giehl. If the conditions are slightly acidic, plants favor the synthesis of the superior Fe^{3+} reductant sideretin, while at high pH, they direct synthesis towards fraxetin, which retains high Fe^{3+} mobilization capacity even under alkaline conditions.

With their work, the researchers provide valuable insights into environment-dependent fine regulation of metabolite biosynthesis and thus help to further understand how [plants](#) adapt to different pH conditions in the soil. The results open up new possibilities for the targeted improvement of plant productivity and plant health under variable soil conditions.

The study is [published](#) in *The Plant Cell* journal.

More information: Vanessa Paffrath et al, A major role of coumarin-dependent ferric iron reduction in strategy I-type iron acquisition in *Arabidopsis*, *The Plant Cell* (2023). [DOI: 10.1093/plcell/koad279](https://doi.org/10.1093/plcell/koad279)

Provided by Leibniz Institute of Plant Genetics and Crop Plant Research

Citation: Plants found to recruit distinct chemical activities of coumarins under different soil pHs (2023, November 29) retrieved 28 April 2024 from <https://phys.org/news/2023-11-distinct-chemical-coumarins-soil-phs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.