

Enhancing the immunosuppressive properties of human umbilical cord mesenchymal stem cells

November 27 2023



Credit: Pixabay/CC0 Public Domain

Mesenchymal stem cells (MSCs) have great potential for the treatment of various immune diseases due to their unique immunomodulatory

properties. However, MSCs exposed to the harsh inflammatory environment of damaged tissue after intravenous transplantation cannot exert their biological effects, and therefore, their therapeutic efficacy is reduced.

In this challenging context, an *in vitro* preconditioning method is necessary for the development of MSC-based therapies with increased immunomodulatory capacity and transplantation efficacy.

To determine whether hypoxia and inflammatory factor preconditioning increase the immunosuppressive properties of MSCs without affecting their biological characteristics, umbilical cord MSCs (UC-MSCs) were pretreated with hypoxia (2% O₂) exposure and inflammatory factors (interleukin-1 β , tumor necrosis factor- α , interferon- γ) for 24 h. The work is [published](#) in the *World Journal of Stem Cells*.

Flow cytometry, [polymerase chain reaction](#), enzyme-linked [immunosorbent assay](#) and other experimental methods were used to evaluate the biological characteristics of pretreated UC-MSCs and to determine whether pretreatment affected the immunosuppressive ability of UC-MSCs in coculture with immune cells.

Pretreatment with hypoxia and inflammatory factors caused UC-MSCs to be elongated but did not affect their viability, proliferation or size. In addition, pretreatment significantly decreased the expression of coagulation-related tissue factors but did not affect the expression of other surface markers. Similarly, mitochondrial function and integrity were retained.

Although pretreatment promoted UC-MSC apoptosis and senescence, it increased the expression of genes and proteins related to immune regulation. Pretreatment increased peripheral blood mononuclear cell and natural killer (NK) cell proliferation rates and inhibited NK cell-

induced toxicity to varying degrees.

In summary, hypoxia and inflammatory factor preconditioning led to higher immunosuppressive effects of MSCs without damaging their biological characteristics.

More information: Hang Li et al, Hypoxia and inflammatory factor preconditioning enhances the immunosuppressive properties of human umbilical cord mesenchymal stem cells, *World Journal of Stem Cells* (2023). [DOI: 10.4252/wjsc.v15.i11.999](https://doi.org/10.4252/wjsc.v15.i11.999)

Provided by World Journal of Stem Cells

Citation: Enhancing the immunosuppressive properties of human umbilical cord mesenchymal stem cells (2023, November 27) retrieved 28 April 2024 from <https://phys.org/news/2023-11-immunosuppressive-properties-human-umbilical-cord.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.