

Oldest known samples of brittle stars from supercontinent Gondwana discovered in South Africa

November 6 2023, by Bob Yirka



Articulated *Krommaster spinosus*. Credit: C. Reddy., CC-BY 4.0 (creativecommons.org/licenses/by/4.0/)

A small team of paleontologists with members from Rhodes University, in South Africa, the National Museum of Natural History, Luxembourg City, and the University of Oxford, in the U.K., has discovered the oldest known brittle star samples from the supercontinent Gondwana at a dig site in South Africa. In their paper [published](#) in the open-access

journal *PLOS ONE*, the group describes where the samples were found, their condition, and where they fit on the ophiuroid family tree.

Brittle stars are often confused with starfish—they both have five [arms](#) attached to a central "body." Brittle stars have much longer arms though (up to 60 cm), which are also much thinner. Also, the arms on most [species](#) of [brittle stars](#) are whip-like, rather than the triangular type found on starfish. Brittle stars use their long arms to grab food as it falls from above or to churn seabed sediment for food.

Brittle stars are echinoderms of the class Ophiuroidea, which is why they are referred to as ophiuroids. They are divided into two main historical categories as well: modern and archaic. Until recently, researchers believed that the archaic species went extinct before the arrival of modern species. But fossil finds (mostly in the [northern hemisphere](#)) have shown that the two co-existed for a time during the Triassic. Thereafter, the archaic species declined in numbers due to an increase in predators during the mid-Paleozoic Marine Revolution.

In this new study, the research team found natural molds of two species of archaic brittle stars in a part of the Bavianskloof Formation, both of which they describe as in near-perfect condition and one of which is a new species—they have named it *Krommaster spinosus*.

The molds were found to be approximately 410 million years old, which makes them the oldest-known brittle star samples found that originated from what was Gondwana, the ancient supercontinent. To better understand their morphology, the research team made silicone replicas from the molds. Their findings may help to better understand the diversification of brittle stars, as almost all other evidence has come from rocks found near the equator.

More information: Caitlin Reddy et al, Earliest known ophiuroids

from high palaeolatitude, southern Gondwana, recovered from the Pragian to earliest Emsian Bavianskloof Formation (Table Mountain Group, Cape Supergroup) South Africa, *PLOS ONE* (2023). [DOI: 10.1371/journal.pone.0292636](https://doi.org/10.1371/journal.pone.0292636)

© 2023 Science X Network

Citation: Oldest known samples of brittle stars from supercontinent Gondwana discovered in South Africa (2023, November 6) retrieved 13 May 2024 from <https://phys.org/news/2023-11-oldest-samples-brittle-stars-supercontinent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.